




ASU College of Integrative
Sciences and Arts
ARIZONA STATE UNIVERSITY



Chiral anomalous effects in magnetars

Igor Shovkovy
Arizona State University

June 21, 2024
University of Science and Technology of China







CHIRAL PLASMA

[Miransky & Shovkovy, Phys. Rep. 576, 1 (2015)]

[Kharzeev, Liao, Voloshin, Wang, Prog. Part. Nucl. Phys. 88, 1 (2016)]

- **Heavy-ion collisions (high temperature)**
[Kharzeev, Liao, Voloshin, Wang, Prog. Part. Nucl. Phys. 88, 1 (2016)]
 - **Super-dense matter in compact stars (high density)**
[Yamamoto, Phys. Rev. D 93, 065017 (2016)]
 - **Early Universe (high temperature)**
[Boyarsky, Frohlich, Ruchayskiy, Phys. Rev. Lett. 108, 031301 (2012)]
 - **Electron plasma in Dirac/Weyl (semi-)metals**
[Gorbar, Miransky, Shovkovy, Sukhachov, *Electronic Properties of Dirac and Weyl Semimetals*, World Scientific, Singapore, 2021]
 - **Other: cold atoms, superfluid $^3\text{He-A}$, etc.**
[Volovik, JETP Lett. 105, 34 (2017)]
- **Magnetospheres of magnetars** [Gorbar & Shovkovy, arXiv:2110.11380]
(electron-positron plasma at moderately high temperature)



- Chiral relativistic plasma may allow $n_L \neq n_R$ to persist on *macroscopic* time/distance scales
- Slow evolution of $n = n_R + n_L$ and $n_5 = n_R - n_L$ is controlled by the continuity equations

$$\frac{\partial n}{\partial t} + \vec{\nabla} \cdot \vec{j} = 0$$

and

$$\frac{\partial n_5}{\partial t} + \vec{\nabla} \cdot \vec{j}_5 = \frac{e^2 \vec{E} \cdot \vec{B}}{2\pi^2} - \Gamma_m n_5$$

where the chirality flip rate: $\Gamma_m \propto \alpha^2 T (m/T)^2$ [Boyarsky et al., PRL **126**, 021801 (2021)]

- Chiral anomaly can produce *macroscopic* effects in plasma




Image credit: NASA

PULSARS


- **Neutron stars** are laboratories of matter under extreme conditions
- Prediction

[Baade & Zwicky, *Proc. Nat. Acad. Sci.* **20**, 259 (1934)]

- Observation

[Hewish, Bell, Pilkington, Scott & Collins, *Nature* **217**, 709 (1968)]

- **Pulsars** are neutron stars that are
 - rapidly rotating ($P \sim 1$ ms to 10 s)
 - strongly magnetized ($B \sim 10^8$ to 10^{15} G)
- Pulsar radiation is beamed along the magnetic field direction (the “lighthouse” effect)



Pulsars in $P-\dot{P}$ plane

- Characteristic age

$$\tau \simeq \frac{P}{2\dot{P}}$$

- Spin-down luminosity

$$-\dot{E} \simeq 4\pi^2 I \frac{\dot{P}}{P^3}$$

- Characteristic magnetic field

$$B \simeq 3 \times 10^{19} \left(\frac{P\dot{P}}{s} \right)^{1/2} \text{ G}$$




Image credit: Condon & Ransom, “Essential Radio Astronomy” (2016)




Image credit: Aurore Simonnet, Sonoma State University

MAGNETOSPHERES

Pulsar electrodynamics (VDM)


- **Vacuum dipole model (VDM)** ($\rho = 0$ & $J = 0$ outside the star)
- Stellar interior (good conductor):

$$\vec{E}'_{in} = \vec{E}_{in} + \frac{\vec{\Omega} \times \vec{r}}{c} \times \vec{B}_{in} = 0$$

- Fields outside the pulsar are

$$\vec{B} = \frac{B_0 R^3}{2r^3} (3(\hat{m} \cdot \hat{r})\hat{r} - \hat{m})$$

$$\vec{E} = \dots \quad [\text{see Deutsch, Ann. Astrophys. 18, 1 (1955)}]$$




where \mathbf{m} is the magnetic moment and $\mathbf{\Omega}$ is the angular frequency

- There is a nonzero charge density and a strong electric field on the surface ($E_{\text{surf}} \sim \Omega R B_0 \sim 10^{12}$ to 10^{15} V/m)

- Charged particles (electrons)
 - i. pulled up from the surface ($\vec{E} \neq 0$)
 - ii. move along curved trajectories ($\vec{B} \neq 0$)
 - iii. produce curvature γ -radiation
 - iv. γ -quanta produce $e^+ e^-$ pairs

$$l_\gamma \simeq \frac{2R_c}{15} \frac{B_c}{B} \frac{m_e}{\epsilon_\gamma}$$

- v. Secondary particles produce synchrotron & curvature radiation
- **Outcome:** Strongly magnetized vacuum is opaque for photons with $\epsilon_\gamma \gtrsim 2m_e$ and, thus, turns into a plasma



- **Rotating magnetosphere model (RMM)** (assuming a highly conducting plasma outside the star)

$$\vec{E}' = \vec{E} + \frac{\vec{\Omega} \times \vec{r}}{c} \times \vec{B} = 0$$

i.e., $E_{\parallel} = 0$


- Plasma motion is determined by

$$\vec{v}_{\text{drift}} = c \frac{\vec{E} \times \vec{B}}{B^2} = \vec{\Omega} \times \vec{r} + j_{\parallel} \vec{B}$$

- Corotating plasma is charged


$$\rho_{\text{GJ}} = \vec{\nabla} \cdot \vec{E} = -\frac{2}{c} \vec{\Omega} \cdot \vec{B}$$

[Goldreich & Julian, *Astrophys. J.* **157**, 869 (1969)]



Gaps in magnetosphere

- If one assumes that $E_{\parallel}=0$ everywhere, the magnetic field lines are equipotential ($V = \text{const}$)



- Then,


$$0 = \oint \vec{E} \cdot d\vec{l} = \int (\vec{\nabla} \times \vec{E}) \cdot d\vec{s} = -\frac{\partial}{\partial t} \int \vec{B} \cdot d\vec{s}$$

- Thus, $E_{\parallel}=0$ cannot be enforced everywhere if \vec{B} changes in time
- Regions (“gaps”) with unscreened E_{\parallel} will necessarily develop (they result from dynamical charge/current starvation)

[Ruderman & Sutherland, *Astrophys. J.* **196**, 51 (1975)]

Gaps in magnetosphere

- Gaps can develop at various locations
- Intermittent gaps are caused by rapid outflow of charge
- The **gap size** h grows at a speed close to the speed of light
- Electric **potential** difference grows like $\Delta V = E_{\parallel} h \propto h^2$
- ΔV & photon flux cause an avalanche production of **electron-positron pairs**
- Since $B \propto 1/r^3$, anomalous effects are strongest near **polar caps**



[Ruderman & Sutherland, *Astrophys. J.* **196**, 51 (1975)]

- Estimates for the **electric field** and the **gap size**

$$\bar{E}_{\parallel} \simeq Bh/R_{LC}$$

$$h \simeq 3.6 \text{ m} \left(\frac{R}{10 \text{ km}} \right)^{2/7} \left(\frac{\Omega}{1 \text{ s}^{-1}} \right)^{-3/7} \left(\frac{B}{10^{14} \text{ G}} \right)^{-4/7}$$

where $R_{LC} = c/\Omega$ is the radius of light cylinder

The field scales with pulsar parameters as follows


$$E_{\parallel} \approx 2.7 \times 10^{-8} E_c \left(\frac{R}{10 \text{ km}} \right)^{2/7} \left(\frac{\Omega}{1 \text{ s}^{-1}} \right)^{4/7} \left(\frac{B}{10^{14} \text{ G}} \right)^{3/7}$$

where $E_c = m_e^2/e = 1.3 \times 10^{18} \text{ V/m}$.

[Ruderman & Sutherland, *Astrophys. J.* **196**, 51 (1975)]

Charge in the gap

formation of the gap



relaxation


$$\eta = \rho / \rho_{GJ}$$

[Timokhin, Mon. Not. R. Astron. Soc. 408, 2092–2114 (2010)]

Charge in gap (simulations)

formation of the gap

formation of plasma



relaxation

$$\eta = \rho / \rho_{GJ}$$

[Timokhin, Mon. Not. R. Astron. Soc. 408, 2092–2114 (2010)]

Gap parameters

- Quantitative estimate of the gap size and fields

B	10^{12} G	10^{13} G	10^{14} G	10^{15} G
h	50 m	13.4 m	3.6 m	0.97 m
$\frac{E_{\parallel}}{E_c}$	3.8×10^{-9}	1.0×10^{-8}	2.7×10^{-8}	7.3×10^{-8}
$\frac{\mathbf{E} \cdot \mathbf{B}}{E_c B_c}$	8.6×10^{-11}	2.3×10^{-9}	6.2×10^{-8}	1.7×10^{-6}

where

$$E_c = m_e^2/e = 1.3 \times 10^{18} \text{ V/m}$$

$$B_c = m_e^2/e = 4.4 \times 10^{13} \text{ G}$$

Chiral charge production

- The evolution of the chiral charge is determined by

$$\frac{\partial n_5}{\partial t} + \vec{\nabla} \cdot \vec{J}_5 = \frac{e^2 \vec{E} \cdot \vec{B}}{2\pi^2} - \Gamma_m n_5$$

- While the chiral anomaly produces n_5 , the chirality flipping tries to wash it away
- The chiral charge n_5 approaches the steady-state value ($t \gg \Gamma_m$):

$$n_5 = \frac{e^2}{2\pi^2 \Gamma_m} \vec{E} \cdot \vec{B}$$

- The estimates for the chirality flip rate in a hot plasma

$$\Gamma_m \simeq \frac{\alpha^2 m_e^2}{T} \quad (T \lesssim m_e / \sqrt{\alpha}) \quad \text{and} \quad \Gamma_m \simeq \frac{\alpha m_e^2}{T} \quad (T \gg m_e / \sqrt{\alpha})$$

[Boyarsky, Cheianov, Ruchayskiy, Sobol, Phys. Rev. Lett. **126**, 021801 (2021)]

- The gap formation time

$$t_h \sim h/c \sim 10^{-8} \text{ s}$$

- Timescale for chiral charge production

$$t^* \sim 1/\Gamma_m \sim 10^{-17} \text{ s}$$

- Note that

$$t_h \gg t^*$$

- Thus, the chirality production is nearly instantaneous

[Gorbar & Shovkovy, Eur. Phys. J. C **82**, 625 (2022)]

Estimate for n_5 in magnetars

- The estimate for the chiral charge is given by

$$n_5 \simeq \frac{e^2 E_{\parallel} B}{2\pi^2 \Gamma_m} \simeq 1.5 \times 10^{-5} \text{ MeV}^3 \left(\frac{T}{1 \text{ MeV}} \right) \\ \times \left(\frac{R}{10 \text{ km}} \right)^{2/7} \left(\frac{\Omega}{1 \text{ s}^{-1}} \right)^{4/7} \left(\frac{B}{10^{14} \text{ G}} \right)^{10/7}$$

- The corresponding chiral chemical potential is

$$\mu_5 \simeq \frac{3n_5}{T^2} \simeq 4.6 \times 10^{-5} \text{ MeV} \left(\frac{T}{1 \text{ MeV}} \right)^{-1} \\ \times \left(\frac{R}{10 \text{ km}} \right)^{2/7} \left(\frac{\Omega}{1 \text{ s}^{-1}} \right)^{4/7} \left(\frac{B}{10^{14} \text{ G}} \right)^{10/7}$$

[Gorbar & Shovkovy, Eur. Phys. J. C **82**, 625 (2022)]

- The corresponding numerical values for chiral charge and chiral chemical potential are

B	10^{12} G	10^{13} G	10^{14} G	10^{15} G
h	50 m	13.4 m	3.6 m	0.97 m
$\frac{E_{\parallel}}{E_c}$	3.8×10^{-9}	1.0×10^{-8}	2.7×10^{-8}	7.3×10^{-8}
$\frac{\mathbf{E} \cdot \mathbf{B}}{E_c B_c}$	8.6×10^{-11}	2.3×10^{-9}	6.2×10^{-8}	1.7×10^{-6}
$\frac{n_5}{m_e^3}$	1.6×10^{-7}	4.3×10^{-6}	1.1×10^{-4}	3.1×10^{-3}
$\frac{\mu_5}{m_e}$	1.2×10^{-7}	3.4×10^{-6}	9.0×10^{-5}	2.4×10^{-3}

[Gorbar & Shovkovy, Eur. Phys. J. C **82**, 625 (2022)]




Image credit: European Southern Observatory


CHIRAL PLASMA INSTABILITY



- Nonzero μ_5 and \vec{B} drive the chiral magnetic effect (CME)

$$\vec{j} = \frac{e^2 \vec{B}}{2\pi^2} \mu_5$$

- The effect comes from the spin-polarized LLL ($s=\downarrow$)
 - **L-handed** states ($p_3 < 0$ & $|E| < \mu_5$) are empty (holes with $p_3 > 0$)
 - **R-handed** states ($p_3 < 0$ & $E < \mu_5$) are occupied



- However, plasma at $\mu_5 \neq 0$ is unstable

- The total current (CME + Ohm)

$$\mathbf{j} = \frac{2\alpha}{\pi} \mu_5 \mathbf{B} + \sigma \mathbf{E}$$

- By substituting \mathbf{j} into Ampere's law

$$\nabla \times \mathbf{B} = \mathbf{j} + \frac{\partial \mathbf{E}}{\partial t}$$

and solving for the electric field, one derives

$$\mathbf{E} = \frac{1}{\sigma} \left(\nabla \times \mathbf{B} - k_\star \mathbf{B} - \frac{\partial \mathbf{E}}{\partial t} \right) \quad \text{where } k_\star = \frac{2\alpha \mu_5}{\pi}$$

- Finally, by calculating the curl and using Faraday's law,

$$\frac{\partial \mathbf{B}}{\partial t} = -\frac{1}{\sigma} \left(\nabla \times (\nabla \times \mathbf{B}) - k_\star \nabla \times \mathbf{B} + \frac{\partial^2 \mathbf{B}}{\partial t^2} \right)$$


Helical modes at $\mu_5 \neq 0$

- Search for a solution as a superposition of helical eigenstates

$$\nabla \times \mathbf{B}_{\lambda,k} = \lambda k \mathbf{B}_{\lambda,k}$$

e.g.,

$$\mathbf{B}_{\lambda,k} = B_0 (\hat{\mathbf{x}} + i\lambda\hat{\mathbf{y}}) e^{-i\omega t + ikz}$$



Then, for a fixed eigenmode, the evolution equation reads

$$\frac{d\mathbf{B}_{\lambda,k}}{dt} = \frac{1}{\sigma} \left(\lambda k_{\star} k - k^2 - \frac{\partial^2}{\partial t^2} \right) \mathbf{B}_{\lambda,k}$$

- The two solutions for the frequency are

$$\omega_{1,2} = -\frac{i}{2} \left(\sigma \pm \sqrt{\sigma^2 + 4k(\lambda k_{\star} - k)} \right)$$

- For a plasma with high conductivity

$$\omega_{1,2} \simeq \begin{cases} -i \left(\sigma + \frac{k(\lambda k_* - k)}{\sigma} \right) \\ i \frac{k(\lambda k_* - k)}{\sigma} \end{cases}$$

[Joyce & Shaposhnikov, PRL 79, 1193 (1997)]
 [Boyarsky, Frohlich, Ruchayskiy, PRL 108, 031301 (2012)]
 [Tashiro, Vachaspati, Vilenkin, PRD 86, 105033 (2012)]
 [Akamatsu & Yamamoto, PRL 111, 052002 (2013)]
 [Tuchin, PRC 91, 064902 (2015)]
 [Manuel & Torres-Rincon, PRD 92, 074018 (2015)]
 [Hirono, Kharzeev, Yin, PRD 92, 125031(2015)]
 [Sigl & Leite, JCAP 01, 025 (2016)]

- The 1st mode is damped by charge screening:

$$B_{k,1} \propto B_0 e^{-\sigma t}$$

- The 2nd mode is unstable when $k < \lambda k_*$:

$$B_{k,2} \propto B_0 e^{+tk(\lambda k_* - k)/\sigma}$$

- The momentum of the fastest growing mode $B_{k,2}$ is

$$\frac{1}{2} k_*$$

Instability in pulsars

- The estimate for k_*

[Gorbar & Shovkovy, Eur. Phys. J. C **82**, 625 (2022)]

$$k_* \simeq 2.2 \times 10^{-7} \text{ MeV} \left(\frac{T}{1 \text{ MeV}} \right)^{-1} \left(\frac{R}{10 \text{ km}} \right)^{2/7} \left(\frac{\Omega}{1 \text{ s}^{-1}} \right)^{4/7} \left(\frac{B}{10^{14} \text{ G}} \right)^{10/7}$$

B	10^{12} G	10^{13} G	10^{14} G	10^{15} G
h	50 m	13.4 m	3.6 m	0.97 m
$\frac{E_{\parallel}}{E_c}$	3.8×10^{-9}	1.0×10^{-8}	2.7×10^{-8}	7.3×10^{-8}
$\frac{\mathbf{E} \cdot \mathbf{B}}{E_c B_c}$	8.6×10^{-11}	2.3×10^{-9}	6.2×10^{-8}	1.7×10^{-6}
$\frac{n_5}{m_e^3}$	1.6×10^{-7}	4.3×10^{-6}	1.1×10^{-4}	3.1×10^{-3}
$\frac{\mu_5}{m_e}$	1.2×10^{-7}	3.4×10^{-6}	9.0×10^{-5}	2.4×10^{-3}
$\frac{k_*}{m_e}$	5.8×10^{-10}	1.6×10^{-8}	4.2×10^{-7}	1.1×10^{-5}

Observational consequences

- Unstable plasma in the gaps produces **helical** (circularly polarized) **modes** in the frequency range

$$0 \lesssim \omega \lesssim k_*$$

- For magnetars, these span **radio frequencies** and may reach into the **near-infrared** range
- Available energy is of the order of $\Delta\mathcal{E} \sim \mu_5^2 T^2 h^3$, i.e.,

$$\Delta\mathcal{E} \simeq 2.1 \times 10^{25} \text{ erg} \left(\frac{T}{1 \text{ MeV}} \right) \left(\frac{R}{10 \text{ km}} \right)^{6/7} \\ \times \left(\frac{\Omega}{1 \text{ s}^{-1}} \right)^{-9/7} \left(\frac{B}{10^{14} \text{ G}} \right)^{2/7}$$

- The energy is sufficient to feed the **fast radio bursts** (?)

Outstanding problems

- Interplay of chiral charge and electron-positron pair **production** induced by energetic photons should be studied in detail
- The modification of the **chiral flip rate** $\Gamma_m \simeq \frac{\alpha^2 m_e^2}{T}$ by the strong magnetic field (extra suppression?)
- The role of the **inverse magnetic cascade** and the **chiral-magnetic turbulence** should be quantified
- Self-consistent **dynamics** of chiral plasma in the gap regions should be simulated in detail
- Detailed mechanism of the **energy transfer** from unstable helical modes to radio emission (FRBs or other types)

[Caleb et al., An emission-state-switching radio transient with a 54-minute period. Nat. Astron. (2024)]

- Chiral anomaly can have *macroscopic* implications in pulsars
- It leads to a *significant* chiral charge production (up to 10^{34} m^{-3}) in strongly magnetized magnetospheres
- The chiral chemical potential μ_5 can be up to 10^{-3} MeV
- This is sufficient to trigger emission of helical waves with frequencies up to about $k_\star \simeq \frac{2}{\pi} \alpha \mu_5$ (radio to infrared range)
- Helical waves can affect the pulsar jets and observable features of the fast radio bursts
- For quantitative effects, further detailed studies are needed