

Generation Function For One-loop Tensor Reduction

Bo Feng^{ab*}

^a*Beijing Computational Science Research Center, Beijing 100084, China*

^b*Peng Huanwu Center for Fundamental Theory, Hefei, Anhui, 230026, China*

ABSTRACT: For loop integrals, the reduction is the standard method. Having an efficient way to find reduction coefficients is an important topic in scattering amplitudes. In this paper, we present the generation functions of reduction coefficients for general one-loop integrals with arbitrary tensor rank.

KEYWORDS: Generation function, Tensor reduction, One-loop integral.

*Emails: fengbo@csrc.ac.cn

Contents

1. Motivation	1
2. Tadpole	3
3. Bubble	4
3.1 Differential equations	5
3.2 The series expansion	7
3.3 The analytic solution	9
3.3.1 The generation function $c_{2\rightarrow 2}$	10
3.3.2 The generation function $c_{2\rightarrow 1;\hat{1}}$	10
4. The general frame	12
4.1 Finding good variables	13
4.2 The differential equations	14
4.3 Analytic solution	15
5. Triangle	19
5.1 The differential equations	20
6. Conclusion	23
A. Solving differential equations	24
B. The explicit solutions of $c_{n,m}$ for bubble reduction	29
B.1 The generation function $c_{2\rightarrow 2}$	29
B.1.1 The generation function $c_{2\rightarrow 1;\hat{1}}$	31
B.1.2 The proof of one useful relation	33

1. Motivation

As the bridge connecting the theoretical frame and the experiment data, scattering amplitude is always one of central concepts in the quantum field theory. Its efficient computation (including higher order contributions in perturbation) becomes the necessary, especially with the coming of the LHC experiment

[1]. The on-shell program¹ in the scattering amplitudes, as the outcome of the challenge, has made the calculations of one-loop amplitude straightforward [10].

In general the loop computation can be divided into two parts: the construction of integrands and then doing the integration. Although the construction of integrands using Feynman diagrams are well established, sometimes it is not the economic way to do so. Looking for a better way to construct integrands is one of current research directions². For the second part, i.e., doing the integration, a very useful method is the reduction. It has been shown that any integral can be written as the linear combination of some basis integrals (called the master integrals) with coefficients as the rational function of external momenta, masses and spacetime dimension. Using the idea of reduction, loop integration can be separated into two parallel tasks: the computation of master integrals and the algorithm to efficiently extract the reduction coefficients. Progresses in any one task will enable us to do more and more complicated integrations (for a nice introduction of recent developments, see [11]).

For the reduction, it can be classified into two categories: the reduction at the integrand level and the reduction at the integral level. The reduction at the integrand level can be systematically solved using the computational algebraic geometry [12, 13, 14, 15]. For the reduction at the integral level, the first proposal is the famous Passarino-Veltman reduction (PV-reduction) method [16]. There are other proposals, such as the Integration-by-Part (IBP) method [17, 18, 19, 20, 21, 22, 23], the unitarity cut method [4, 5, 7, 24, 25, 26, 27, 28, 29, 30], Intersection number [31, 32, 33, 34]. Although there are a lot of developments for the reduction at the integral level, it is still desirable to improve them by current complexity of computations.

In recent papers [35, 36, 37, 39, 38] we have introduced the auxiliary vector R to improve the traditional PV-reduction method. Using R we can construct the differential operators and then establish algebraic recurrence relation to determine reduction coefficients analytically. This method has also been generalized to two-loop sunset diagram (see [38]) where the original PV-reduction method is hard to be used. When using the auxiliary vector R in the IBP method, the efficiency of reduction has also been improved as shown in [40, 41].

Although the advantage of using auxiliary vector R has been demonstrated from various aspects, the algebraic recursive structure makes it still hard to have a general understanding of reduction coefficients for higher and higher tensor ranks. Could we get more by this method? As we will show in this paper, indeed we can get more if we probe the reduction problem from a new angle. The key idea is the concept of **generation function**. In fact, generation function is well known in physics and mathematics. Sometimes the coefficients of a series is hard to imagine, but the series itself is easy to write down. For example, the

¹There are many works in this topic. For introduction, please see following two books [2, 3]. Some early works with on-shell concept are [4, 5, 6, 7, 8, 9].

²For example, the unitarity cut method proposed in [4, 5, 7] uses on-shell tree level amplitudes as input.

Hermite Polynomial $H_n(x)$ can be read out from the generation function

$$e^{2tx-t^2} = \sum_{n=0}^{\infty} H_n(x) \frac{t^n}{n!}. \quad (1.1)$$

Thus we can ask that if we sum the reduction coefficients of different tensor ranks together, could we get a simpler answer? For reduction problem, the numerator of tensor rank k is given by $(2\ell \cdot R)^k$ in our method and we need to see how to sum them together. There are many ways to sum them. Two typical ones are

$$\psi_1(t) = \sum_{n=0}^{\infty} t^n (2\ell \cdot R)^n = \frac{1}{1 - t(2\ell \cdot R)}, \quad \psi_2(t) = \sum_{n=0}^{\infty} \frac{(2\ell \cdot R)^n t^n}{n!} = e^{t(2\ell \cdot R)}. \quad (1.2)$$

In this paper, we will focus on the generation function of the type $\psi_2(t)$ because it is invariant under the differential action, i.e, $\frac{de^x}{dx} = e^x$. We will see that the generation functions satisfy simple differential equations, which can be solved analytically.

The plan of the paper is following. In the section two, we present the generation function of reduction coefficients of tadpole integral. The tadpole example is a little bit trivial, thus in the section three, we discuss carefully how to find generation functions for the bubble integral, which is the simplest nontrivial example. With the experience obtained for bubble, we present the solution for general one-loop integrals in the section four. To demonstrate the frame established in section four, we discuss briefly the triangle example in the section five. Finally, a brief summary and discussion is given in the section six. Some technical details have been collected in the Appendix, where in the Appendix A, the solution of two typical differential equations is presented, while the solution of recursive relation for the bubble has been explained in the Appendix B.

2. Tadpole

With above brief discussion, in this section we start from the simplest case, i.e., the tadpole topology to discuss the generation function. Summing all tensor ranks properly we have³

$$I_{tad}(t, R) \equiv \int d\ell \frac{e^{t(2\ell \cdot R)}}{\ell^2 - M^2} = c_{1 \rightarrow 1}(t, R, M) \int d\ell \frac{1}{\ell^2 - M^2} \quad (2.1)$$

where $c_{1 \rightarrow 1}(t, R, M)$ is the generation function of reduction coefficients and for simplicity we have defined $\int d\ell_i(\bullet) \equiv \int \frac{d^D \ell_i}{i\pi^{D/2}}(\bullet)$. To find closed analytic expression for $c_{1 \rightarrow 1}(t, R, M)$ we establish the corresponding differential equation. Acting with ∂_R we have

$$\frac{\partial}{\partial R} \cdot \frac{\partial}{\partial R} I_{tad}(t, R) = \int d\ell \frac{4t^2 \ell^2 e^{t(2\ell \cdot R)}}{\ell^2 - M^2} = 4t^2 M^2 I_{tad}(t, R) \quad (2.2)$$

³The mass dimension of parameter t is -2 .

at one side, and $(\frac{\partial}{\partial R} \cdot \frac{\partial}{\partial R} c_{1 \rightarrow 1}(t, R, M)) \int d\ell \frac{1}{\ell^2 - M^2}$ at another side, thus we get

$$\frac{\partial}{\partial R} \cdot \frac{\partial}{\partial R} c_{1 \rightarrow 1}(t, R, M) = 4t^2 M^2 c_{1 \rightarrow 1}(t, R, M) . \quad (2.3)$$

By the Lorentz invariance, $c_{1 \rightarrow 1}(t, R, M)$ is the function of $r = R \cdot R$ only, i.e., $c_{1 \rightarrow 1}(t, R, M) = f(r)$. It is easy to see that differential equation (2.3) becomes

$$\boxed{4r f'' + 2D f' - 4t^2 M^2 f = 0} , \quad (2.4)$$

which is the form (A.14) studied in the Appendix A. This second order differential equation has two singular points $r = 0$ and $r = \infty$, where the singular point $r = 0$ is canonical. The solution has been given in (A.33). Putting $A = 4, B = 2D, C = -4t^2 M^2$ in (A.29) and the boundary condition $c_0 = 1$, we get immediately

$$c_{1 \rightarrow 1}(t, R, M) = \sum_{n=0}^{\infty} \frac{(t^2 M^2 r)^n}{n! (\frac{D}{2})_n} = {}_0F_1(\emptyset; \frac{D}{2}; t^2 M^2 r) . \quad (2.5)$$

Before ending this section, let us mention that when we do the reduction for other topologies, we will meet the reduction of $\int d\ell \frac{e^{t(2\ell \cdot R)}}{(\ell - K)^2 - M^2}$. Using the momentum shifting, it is easy to see that

$$\int d\ell \frac{e^{t(2\ell \cdot R)}}{(\ell - K)^2 - M^2} = \int d\ell \frac{e^{t(2(\ell+K) \cdot R)}}{\ell^2 - M^2} = e^{2t(K \cdot R)} c_{1 \rightarrow 1}(t, R, M) \int d\ell \frac{1}{\ell^2 - M^2} . \quad (2.6)$$

The results shows the advantage of using generation function with exponential form.

3. Bubble

Having found the generation function of tadpole reduction, we move to the first nontrivial example, i.e., the generation function of bubble reduction, which is defined through

$$\begin{aligned} I_{bub}(t, R) &\equiv \int d\ell \frac{e^{t(2\ell \cdot R)}}{(\ell^2 - M_0^2)((\ell - K)^2 - M_1^2)} \\ &= c_{2 \rightarrow 2} \int d\ell \frac{1}{(\ell^2 - M_0^2)((\ell - K)^2 - M_1^2)} + c_{2 \rightarrow 1; \hat{1}} \int d\ell \frac{1}{(\ell^2 - M_0^2)} + c_{2 \rightarrow 1; \hat{0}} \int d\ell \frac{1}{((\ell - K)^2 - M_1^2)} . \end{aligned} \quad (3.1)$$

For simplicity, we have not written down variables of reduction coefficients explicitly. If written explicitly, it will be $c(t, R, K; M_0, M_1)$ or $c(t, R^2, K \cdot R, K^2; M_0, M_1)$ if using the Lorentz contraction form.

The generation form (3.1) can produce some nontrivial relations among these generation functions of

reduction coefficients easily. Noticing that⁴

$$\begin{aligned}
I_{bub}(t, R) &\equiv \int d\ell \frac{e^{t(2\ell \cdot R)}}{(\ell^2 - M_0^2)((\ell - K)^2 - M_1^2)} = e^{t(2K \cdot R)} \int d\tilde{\ell} \frac{e^{t(2\tilde{\ell} \cdot R)}}{((\tilde{\ell} + K)^2 - M_0^2)(\tilde{\ell}^2 - M_1^2)} \\
&= e^{t(2K \cdot R)} \left\{ c_{2 \rightarrow 2}(t, R^2, -K \cdot R, K^2; M_1, M_0) \int d\ell \frac{1}{(\ell^2 - M_0^2)((\ell - K)^2 - M_1^2)} \right. \\
&\quad + c_{2 \rightarrow 1; \hat{0}}(t, R^2, -K \cdot R, K^2; M_1, M_0) \int d\ell \frac{1}{(\ell^2 - M_0^2)} \\
&\quad \left. + c_{2 \rightarrow 1; \hat{1}}(t, R^2, -K \cdot R, K^2; M_1, M_0) \int d\ell \frac{1}{((\ell - K)^2 - M_1^2)} \right\} , \tag{3.2}
\end{aligned}$$

we have

$$c_{2 \rightarrow 2}(t, R^2, K \cdot R, K^2; M_0, M_1) = e^{t(2K \cdot R)} c_{2 \rightarrow 2}(t, R^2, -K \cdot R, K^2; M_1, M_0) , \tag{3.3}$$

$$c_{2 \rightarrow 1; \hat{1}}(t, R^2, K \cdot R, K^2; M_0, M_1) = e^{t(2K \cdot R)} c_{2 \rightarrow 1; \hat{0}}(t, R^2, -K \cdot R, K^2; M_1, M_0) , \tag{3.4}$$

$$c_{2 \rightarrow 1; \hat{0}}(t, R^2, K \cdot R, K^2; M_0, M_1) = e^{t(2K \cdot R)} c_{2 \rightarrow 1; \hat{1}}(t, R^2, -K \cdot R, K^2; M_1, M_0) \tag{3.5}$$

by comparing (3.2) with (3.1). The first relation (3.3) can be a consistent check for $c_{2 \rightarrow 2}$ while the second relation (3.4) and the third relation (3.5) tell us that we need to compute only one of $c_{2 \rightarrow 1; \hat{i}}$ functions. Another useful check is the mass dimension. Since the mass dimension of t is (-2) , we have

$$[c_{2 \rightarrow 2}] = 0, \quad [c_{2 \rightarrow 1}] = -2 \tag{3.6}$$

3.1 Differential equations

Now we will write down differential equations for these generation functions. Acting $\partial_R \cdot \partial_R$ on both sides of (3.1) we have

$$\begin{aligned}
\partial_R \cdot \partial_R I_{bub}(t, R) &= \int d\ell \frac{4t^2 \ell^2 e^{t(2\ell \cdot R)}}{(\ell^2 - M_0^2)((\ell - K)^2 - M_1^2)} \\
&= 4t^2 M_0^2 I_{bub}(t, R) + 4t^2 e^{t(2K \cdot R)} \int d\ell \frac{e^{t(2\ell \cdot R)}}{(\ell^2 - M_1^2)} , \tag{3.7}
\end{aligned}$$

⁴Comparing to the shifting symmetry discussed in (3.2), one can also consider the symmetry with $\ell \rightarrow -\ell$. For this one, we have $R \rightarrow -R$ and $K \rightarrow -K$. If using the variables $R^2, K \cdot R$, it is invariant. In other words, the symmetry $\ell \rightarrow -\ell$ is trivial.

thus we derive

$$\partial_R \cdot \partial_R c_{2 \rightarrow 2}(t, R^2, K \cdot R, K^2; M_0, M_1) = 4t^2 M_0^2 c_{2 \rightarrow 2}(t, R^2, K \cdot R, K^2; M_0, M_1) , \quad (3.8)$$

$$\partial_R \cdot \partial_R c_{2 \rightarrow 1; \widehat{1}}(t, R^2, K \cdot R, K^2; M_0, M_1) = 4t^2 M_0^2 c_{2 \rightarrow 1; \widehat{1}}(t, R^2, K \cdot R, K^2; M_0, M_1) , \quad (3.9)$$

$$\begin{aligned} \partial_R \cdot \partial_R c_{2 \rightarrow 1; \widehat{0}}(t, R^2, K \cdot R, K^2; M_0, M_1) &= 4t^2 M_0^2 c_{2 \rightarrow 1; \widehat{0}}(t, R^2, K \cdot R, K^2; M_0, M_1) \\ &+ 4t^2 e^{t(2K \cdot R)} c_{1 \rightarrow 1}(t, R^2, M_1) . \end{aligned} \quad (3.10)$$

Acting with $K \cdot \partial_R$ we have

$$\begin{aligned} K \cdot \partial_R I_{bub}(t, R) &= \int d\ell \frac{t(2K \cdot \ell) e^{t(2\ell \cdot R)}}{(\ell^2 - M_0^2)((\ell - K)^2 - M_1^2)} = \int d\ell \frac{t(D_0 - D_1 + f) e^{t(2\ell \cdot R)}}{(\ell^2 - M_0^2)((\ell - K)^2 - M_1^2)} \\ &= t f I_{bub}(t, R) - t \int d\ell \frac{e^{t(2\ell \cdot R)}}{(\ell^2 - M_0^2)} + t e^{t(2K \cdot R)} \int d\ell \frac{e^{t(2\ell \cdot R)}}{(\ell^2 - M_1^2)} , \end{aligned} \quad (3.11)$$

thus we derive

$$K \cdot \partial_R c_{2 \rightarrow 2}(t, R^2, K \cdot R, K^2; M_0, M_1) = t f c_{2 \rightarrow 2}(t, R^2, K \cdot R, K^2; M_0, M_1) , \quad (3.12)$$

$$K \cdot \partial_R c_{2 \rightarrow 1; \widehat{1}}(t, R^2, K \cdot R, K^2; M_0, M_1) = t f c_{2 \rightarrow 1; \widehat{1}}(t, R^2, K \cdot R, K^2; M_0, M_1) - t c_{1 \rightarrow 1}(t, R^2, M_0) , \quad (3.13)$$

$$K \cdot \partial_R c_{2 \rightarrow 1; \widehat{0}}(t, R^2, K \cdot R, K^2; M_0, M_1) = t f c_{2 \rightarrow 1; \widehat{0}}(t, R^2, K \cdot R, K^2; M_0, M_1) + t e^{t(2K \cdot R)} c_{1 \rightarrow 1}(t, R^2, M_1) \quad (3.14)$$

where $f = K^2 - M_1^2 + M_0^2$. Above two groups of differential equations can be uniformly written as

$$\partial_R \cdot \partial_R c_T = 4t^2 M_0^2 c_T + 4t^2 \xi_R h_T , \quad (3.15)$$

$$K \cdot \partial_R c_T = t f c_T + t \xi_K h_T \quad (3.16)$$

where h_T is the possible non-homogenous contribution coming from lower topology (tadpole). For different T we have

$$T = \{2 \rightarrow 2\} : \quad h_T = 0 \quad \text{or} \quad \xi_R = \xi_K = 0 , \quad (3.17)$$

$$T = \{2 \rightarrow 1; \widehat{1}\} : \quad h_T = c_{1 \rightarrow 1}(t, R^2, M_0), \quad \xi_R = 0, \quad \xi_K = -1 , \quad (3.18)$$

$$T = \{2 \rightarrow 1; \widehat{0}\} : \quad h_T = e^{t(2K \cdot R)} c_{1 \rightarrow 1}(t, R^2, M_1), \quad \xi_R = 1, \quad \xi_K = +1 . \quad (3.19)$$

Generation functions are functions of $f(r, p)$ with $r = R^2$ and $p = K \cdot R$. It is easy to work out that

$$\frac{\partial}{\partial R^\mu} = 2\eta_{\rho\mu} R^\rho \partial_r + K_\mu \partial_p, \quad K \cdot \frac{\partial}{\partial R^\mu} = 2p \partial_r + K^2 \partial_p \quad (3.20)$$

$$\eta^{\mu\nu} \frac{\partial}{\partial R^\nu} \frac{\partial}{\partial R^\mu} = (4r \partial_r^2 + 4p \partial_p \partial_r + K^2 \partial_p^2 + 2D \partial_r) \quad (3.21)$$

thus (3.15) and (3.16) can be written as

$$(4r\partial_r^2 + 4p\partial_p\partial_r + K^2\partial_p^2 + 2D\partial_r - 4t^2M_0^2)c_T = 4t^2\xi_R h_T \quad , \quad (3.22)$$

$$(2p\partial_r + K^2\partial_p - tf)c_T = t\xi_K h_T \quad . \quad (3.23)$$

Equations (3.22) and (3.23) are the differential equations we need to solve. We will present two ways to solve them. One is by the series expansion of the naive variables r, p . This is the method used in [35, 36]. However, as we will show, using the idea of generation function, the powers of r, p are independent to each other, thus the recursion relations become simpler and can be solved explicitly. Another method is to solve the differential equation directly and get more compact and analytical expression. An important lesson from the second method is that the right variables to do the series expansion are not r, p but their proper combination.

3.2 The series expansion

In this subsection, we will present the series expansion of r, p . Writing⁵

$$c = \sum_{n,m=0}^{\infty} c_{n,m} r^n p^m, \quad h_c = \sum_{n,m=0}^{\infty} h_{n,m} r^n p^m \quad (3.24)$$

and putting them to (3.22) and (3.23) we get following equations

$$0 = 2(n+1)(2n+2m+D)c_{n+1,m} + K^2(m+2)(m+1)c_{n,m+2} - 4t^2M_0^2c_{n,m} - 4t^2\xi_R h_{n,m} \quad n, m \geq 0 \quad (3.25)$$

$$0 = 2(n+1)c_{n+1,m} + K^2(m+2)c_{n,m+2} - tfc_{n,m+1} - t\xi_K h_{n,m+1} \quad m, n \geq 0 \quad (3.26)$$

$$0 = K^2c_{n,1} - tfc_{n,0} - t\xi_K h_{n,0} \quad n \geq 0 \quad (3.27)$$

Using (3.25) and (3.26) we can solve

$$\begin{aligned} c_{n+1,m} &= \frac{4M_0^2t^2}{2(n+1)(D+2n+m-1)}c_{n,m} - \frac{tf(m+1)}{2(n+1)(D+2n+m-1)}c_{n,m+1} \\ &+ \frac{4t^2\xi_R h_{n,m} - t\xi_K(m+1)h_{n,m+1}}{2(n+1)(D+2n+m-1)} \end{aligned} \quad (3.28)$$

$$\begin{aligned} c_{n,m+2} &= \frac{-4M_0^2t^2}{(m+2)(D+2n+m-1)K^2}c_{n,m} + \frac{tf(D+2n+2m)}{(m+2)(D+2n+m-1)K^2}c_{n,m+1} \\ &+ \frac{-4t^2\xi_R h_{n,m} + t\xi_K(D+2n+2m)h_{n,m+1}}{2(n+1)(D+2n+m-1)} \end{aligned} \quad (3.29)$$

⁵Since we are considering the generation functions, the n, m are free indices, while in previous works [35, 36] with fixed tensor rank K , n, m are constrained by $2n+m=K$. One can see that many manipulations are simplified using the idea of generation functions.

Now setting $m = 0$ in (3.28) and combining (3.27) we can solve

$$c_{n+1,0} = \frac{(-t^2 f^2 + 4M_0^2 t^2 K^2)}{2(n+1)(D+2n-1)K^2} c_{n,0} - \frac{t^2(\xi_K f - 4\xi_R K^2)h_{n,0} + t\xi_K K^2 h_{n,1}}{2(n+1)(D+2n-1)K^2} \quad (3.30)$$

$$c_{n,1} = \frac{tf}{K^2} c_{n,0} + \frac{t\xi_K h_{n,0}}{K^2} \quad (3.31)$$

Using equation (3.30) we can recursively solve all $c_{n,0}$ starting from the boundary condition $c_{0,0} = 1$ for $c_{2 \rightarrow 2}$ or $c_{0,0} = 0$ for $c_{2 \rightarrow 1}$. Knowing all $c_{n,0}$ we can use (3.31) to get all $c_{n,1}$. To solve all $c_{n,m}$, we use (3.25) and (3.26) again, but now solve

$$c_{n,m+1} = \frac{4M_0^2 t}{f(m+1)} c_{n,m} - \frac{2(n+1)(D+2n+m-1)}{tf(m+1)} c_{n+1,m} + 4t\xi_R h_{n,m} - \xi_K(m+1)h_{n,m+1} \quad (3.32)$$

$$c_{n,m+2} = \frac{-2(n+1)(D+2n+2m)}{(m+1)(m+2)K^2} c_{n+1,m} + \frac{4t^2 M_0^2}{(m+1)(m+2)K^2} c_{n,m} + \frac{4t^2 \xi_R h_{n,m}}{(m+1)(m+2)K^2} \quad (3.33)$$

Both equations can be used recursively to solve $c_{n,m}$. After using one of them to get all $c_{n,m}$, another one becomes nontrivial consistent check. Among them (3.32) is better, since it solves $(m+1)$ from m .

Using above algorithm, we present the first few terms of generation functions for comparison. For $c_{2 \rightarrow 2}$ we have

$$\begin{aligned} c_{2,2} = & 1 + \frac{ft}{K^2 p} + \frac{(Df^2 - 4K^2 M_0^2)t^2}{2(D-1)(K^2)^2} p^2 + \frac{(4K^2 M_0^2 - f^2)t^2}{2(D-1)K^2} r \\ & + \frac{f((2+D)f^2 - 12K^2 M_0^2)t^3}{6(D-1)(K^2)^3} p^3 - \frac{f(f^2 - 4K^2 M_0^2)t^3}{2(D-1)(K^2)^2} r p + \dots \end{aligned} \quad (3.34)$$

and

$$\begin{aligned} c_{2 \rightarrow 1; \hat{1}} = & 0 - \frac{t}{K^2 p} - \frac{Dft^2}{2(D-1)(K^2)^2} p^2 + \frac{ft^2}{2(D-1)K^2} r \\ & - \frac{(D(2+D)f^2 - 8(D-1)K^2 M_0^2)t^3}{6(D-1)D(K^2)^3} p^3 + \frac{(Df^2 - 4(D-1)K^2 M_0^2)t^3}{2(D-1)D(K^2)^2} r p + \dots \end{aligned} \quad (3.35)$$

and

$$\begin{aligned} c_{2 \rightarrow 1; \hat{0}} = & 0 + \frac{t}{K^2 p} + \frac{(-D\tilde{f} + 4(D-1)K^2)t^2}{2(D-1)(K^2)^2} p^2 + \frac{\tilde{f}t^2}{2(D-1)K^2} r + \frac{(Df\tilde{f} + 4(D-1)K^2 M_1^2)t^3}{2(D-1)D(K^2)^2} r p \\ & + \frac{(6DK^2(K^2(D-2) + DM_0^2) - 2(D+2)(3D-2)K^2 M_1^2 + D(D+2)\tilde{f}^2)t^3}{6(D-1)(K^2)^2} p^3 + \dots \end{aligned} \quad (3.36)$$

where we have defined $\tilde{f} = K^2 - M_0^2 + M_1^2 = -f + 2K^2$.

Here we have presented the general recursive algorithm. In the Appendix B, we will show that these

recursion relations can be solved explicitly, i.e., we find explicit expressions for all coefficients $c_{n,m}$.

3.3 The analytic solution

In previous subsection, we have present the solution using the series expansion. In this subsection, we will solve the two differential equations (3.22) and (3.23) directly.

Let us start from (3.23) first. To solve it, we define following new variables

$$x = K^2 r - p^2, \quad y = p, \quad (3.37)$$

then (3.23) becomes

$$(2p\partial_r + K^2\partial_p - tf)c(r, p) = (K^2\partial_y - tf)c(x, y) = t\xi_K h_T, \quad (3.38)$$

where we have used

$$p = y, \quad r = \frac{x + y^2}{K^2}, \quad \partial_r = K^2\partial_x, \quad \partial_p = -2y\partial_x + \partial_y. \quad (3.39)$$

The differential equation (3.38) can be solved as (see the discussion in the Appendix, for example, (A.5))

$$c(x, y) = \frac{1}{K^2} e^{\frac{tf}{K^2}y} \left(G(x) + \int_0^y dw e^{-\frac{tf}{K^2}w} t\xi_K h_T(x, w) \right), \quad (3.40)$$

where the function G depends only on x , while $h_T \equiv h_T(x, y)$ is the function of both x, y .

Now we consider the equation (3.22). The first step is to simplify it by writing

$$(4r\partial_r^2 + 4p\partial_p\partial_r + K^2\partial_p^2 + 2D\partial_r) = \frac{1}{K^2} (2p\partial_r + K^2\partial_p)(2p\partial_r + K^2\partial_p) + (4r - \frac{4p^2}{K^2})\partial_r^2 + 2(D-1)\partial_r. \quad (3.41)$$

Thus using (3.23), (3.22) becomes

$$\left(4xK^2\partial_x^2 + 2(D-1)K^2\partial_x + \frac{t^2(f^2 - 4K^2M_0^2)}{K^2} \right) c = \left(-t\xi_K\partial_y + \frac{4t^2\xi_R K^2 - t^2\xi_K f}{K^2} \right) h_T. \quad (3.42)$$

Putting (3.40) to (3.42) and simplifying we get

$$\begin{aligned} & \left(4xK^2\partial_x^2 + 2(D-1)K^2\partial_x + \frac{t^2(f^2 - 4K^2M_0^2)}{K^2} \right) G(x) \\ &= K^2 e^{-\frac{tf}{K^2}y} \left(-t\xi_K\partial_y + \frac{4t^2\xi_R K^2 - t^2\xi_K f}{K^2} \right) h_T(x, y) \\ & - \left(4xK^2\partial_x^2 + 2(D-1)K^2\partial_x + \frac{t^2(f^2 - 4K^2M_0^2)}{K^2} \right) \int_0^y dw e^{-\frac{tf}{K^2}w} t\xi_K h_T(x, w). \end{aligned} \quad (3.43)$$

Equation (3.43) is the form of (A.14) which has been discussed in the Appendix. One interesting point is

that since the left hand side is independent of y , the right hand side should be zero under the action of ∂_y . One can check that it is indeed true.

Having laid out the frame, we can use it to solve various generation functions.

3.3.1 The generation function $c_{2 \rightarrow 2}$

For this case, we have $h_T = 0$, thus using the result (A.29) we can immediately written down

$$c_{2 \rightarrow 2}(t, r, p, K^2; M_0, M_1) = {}_0F_1(\emptyset; \frac{D-1}{2}; \left(\frac{t^2(4K^2M_0^2 - f^2)x}{4(K^2)^2} \right) e^{\frac{tf}{K^2}y} |_{x \rightarrow K^2r - p^2, y \rightarrow p} \quad (3.44)$$

One can check it with the series expansion (B.3) given in the Appendix B. Comparing to it, the result (3.44) is very simple and compact. This shows the power of using the generation function. Also, the differential equations (3.22) and (3.23) tell us the right variables for the series expansion should be x, y instead of the naive variables r, p .

3.3.2 The generation function $c_{2 \rightarrow 1; \hat{1}}$

For this case we have $\xi_R = 0, \xi_K = -1$ and

$$h_T(r) = c_{1 \rightarrow 1}(t, r = \frac{x + y^2}{K^2}, M_0) = \sum_{n=0}^{\infty} \frac{(t^2 M_0^2)^n \left(\frac{x+y^2}{K^2} \right)^n}{n! \left(\frac{D}{2} \right)_n} \quad (3.45)$$

which satisfies the differential equation (2.4). The (3.43) becomes

$$\begin{aligned} & \left(4xK^2\partial_x^2 + 2(D-1)K^2\partial_x + \frac{t^2(f^2 - 4K^2M_0^2)}{K^2} \right) G(x) \\ &= K^2 e^{-\frac{tf}{K^2}y} t \left(\partial_y + \frac{tf}{K^2} \right) h_T(x, y) \\ &+ t \left(4xK^2\partial_x^2 + 2(D-1)K^2\partial_x + \frac{t^2(f^2 - 4K^2M_0^2)}{K^2} \right) \int_0^y dw e^{-\frac{tf}{K^2}w} h_T(x, w) \end{aligned} \quad (3.46)$$

A first important check is that the right hand side of (3.46) is y -independent. Acting $\frac{\partial}{\partial y}$ at the right hand side we will get

$$\begin{aligned} & t e^{-\frac{tf}{K^2}y} \left\{ -tf \left(\partial_y + \frac{tf}{K^2} \right) h_T(x, y) + K^2 \left(\partial_y^2 + \frac{tf}{K^2} \partial_y \right) h_T(x, y) \right. \\ & \left. + \left(4xK^2\partial_x^2 + 2(D-1)K^2\partial_x + \frac{t^2(f^2 - 4K^2M_0^2)}{K^2} \right) h_T(x, y) \right\} \end{aligned} \quad (3.47)$$

Using

$$\partial_x h_T(r) = \frac{\partial \frac{x+y^2}{K^2}}{\partial x} \partial_r h_T = \frac{1}{K^2} \partial_r h_T, \quad \partial_y h_T(r) = \frac{\partial \frac{x+y^2}{K^2}}{\partial y} \partial_r h_T = \frac{2y}{K^2} \partial_r h_T \quad (3.48)$$

one can check that (3.47) is reduced to the differential equation (2.4), thus we have proved the y -independent of (3.46).

Setting $y = 0$ in (3.46) we get

$$\left(4xK^2 \partial_x^2 + 2(D-1)K^2 \partial_x + \frac{t^2(f^2 - 4K^2 M_0^2)}{K^2} \right) G(x) = t^2 f h_T(x, y=0) \quad (3.49)$$

where we have used $\partial_y h_T(x, y=0) = \frac{2y}{K^2} \partial_r h_T = 0$. The differential equation (3.49) is the form of (A.14) and we get the solution

$$G(x) = \frac{t^2 f}{4K^2} G_0(x) \int_0^x dw w^{-\frac{(D-1)}{2}} e^{-2G_0(w)} \int_0^w d\xi h_T(\xi, y=0) G_0^{-1}(\xi) e^{2G_0(\xi)} \xi^{\frac{(D-1)}{2}-1} \quad (3.50)$$

where

$$G_0(x) = {}_0F_1\left(\emptyset; \frac{(D-1)}{2}; \frac{t^2(4K^2 M_0^2 - f^2)}{4(K^2)^2} x\right) \quad (3.51)$$

Putting all together we finally have

$$c_{2 \rightarrow 1; \hat{1}}(x, y) = \frac{1}{K^2} e^{\frac{tf}{K^2} y} \left(G(x) - t \int_0^y dw e^{-\frac{tf}{K^2} w} h_T(x, w) \right) \quad (3.52)$$

Although we have a very compact expression (3.52) for generation function, in practice it is more desirable to have the series expansion form. In the Appendix A we have introduced three ways. Here we work out the expansion by direct integration. Using (3.45) we have

$$\int_0^y dw e^{-\frac{tf}{K^2} w} h_T(x, w) = \sum_{n=0}^{\infty} \frac{(t^2 M_0^2)^n}{n! \left(\frac{D}{2}\right)_n} \int_0^y dw e^{-\frac{tf}{K^2} w} \left(\frac{x+w^2}{K^2} \right)^n \quad (3.53)$$

To work out the integration, we see that

$$R(\alpha) \equiv \int_0^T du e^{\alpha u} = \frac{1}{\alpha} e^{\alpha u} \Big|_0^T = \frac{e^{\alpha T} - 1}{\alpha} \quad (3.54)$$

thus

$$\begin{aligned} \frac{d^n}{d\alpha^n} R(\alpha) &= \int_0^T du e^{\alpha u} u^n = \frac{(-)^n n!}{\alpha^{n+1}} (e^{\alpha T} [e^{-\alpha T}]_{\alpha^n} - 1) \\ &= \frac{(-)^n n!}{\alpha^{n+1}} \left(e^{\alpha T} \left(e^{-\alpha T} - \sum_{i=n+1}^{\infty} \frac{(-\alpha T)^i}{i!} \right) - 1 \right) = \frac{-(-)^n n!}{\alpha^{n+1}} e^{\alpha T} \sum_{i=n+1}^{\infty} \frac{(-\alpha T)^i}{i!} \end{aligned} \quad (3.55)$$

where the symbol $[Y(x)]_{x^{n-1}}$ means to keep the Taylor expansion up to the order of $x^{(n-1)}$. Using (3.55) we have

$$\begin{aligned} \int_0^T du e^{\alpha u} (\beta + \gamma u^2)^N &= \sum_{i=0}^N \frac{N!}{i!(N-i)!} \beta^{N-i} \gamma^i \int_0^T du e^{\alpha u} u^{2i} \\ &= \sum_{i=0}^N \frac{N!}{i!(N-i)!} \beta^{N-i} \gamma^i \frac{(2i)!}{\alpha^{2i+1}} e^{\alpha T} \sum_{j=2i+1}^{\infty} \frac{(-\alpha T)^j}{j!} \end{aligned} \quad (3.56)$$

Using (3.56) we can evaluate (3.53) as

$$\int_0^y dw e^{-\frac{tf}{K^2} w} h_T(x, w) = y e^{-\frac{tf}{K^2} y} \sum_{n=0}^{\infty} \sum_{j=0}^{\infty} \sum_{i=0}^n \frac{\left(\frac{t^2 M_0^2}{K^2}\right)^n}{\left(\frac{D}{2}\right)_n} \frac{(2i)! x^{n-i} y^{2i}}{i!(n-i)!} \frac{\left(\frac{tfy}{K^2}\right)^j}{(j+2i+1)!} \quad (3.57)$$

The evaluation of $G(x)$ can be found in (A.23) as

$$G(x) = \sum_{n=0}^{\infty} \sum_{i=0}^{n-1} \frac{\left(\frac{D-1}{2}\right)_i}{n! \left(\frac{D-1}{2}\right)_n \left(\frac{D}{2}\right)_i} \frac{f(M_0^2)^i (t^2 x)^n (4K^2 M_0^2 - f^2)^{n-i-1}}{4^{n-i} (K^2)^{2n-i-1}} \quad (3.58)$$

Collecting all pieces together, we finally have

$$\begin{aligned} c_{2 \rightarrow 1; \hat{1}}(t, r, p, K^2; M_0, M_1) &= \frac{1}{K^2} e^{\frac{tf}{K^2} y} \sum_{n=0}^{\infty} \sum_{i=0}^{n-1} \frac{\left(\frac{D-1}{2}\right)_i}{n! \left(\frac{D-1}{2}\right)_n \left(\frac{D}{2}\right)_i} \frac{f(M_0^2)^i (t^2 x)^n (4K^2 M_0^2 - f^2)^{n-i-1}}{4^{n-i} (K^2)^{2n-i-1}} \\ &\quad - \frac{ty}{K^2} \sum_{n=0}^{\infty} \sum_{j=0}^{\infty} \sum_{i=0}^n \frac{\left(\frac{t^2 M_0^2}{K^2}\right)^n}{\left(\frac{D}{2}\right)_n} \frac{(2i)! x^{n-i} y^{2i}}{i!(n-i)!} \frac{\left(\frac{tfy}{K^2}\right)^j}{(j+2i+1)!} \end{aligned} \quad (3.59)$$

This result can be checked with the one given in (B.14). One can see that the formula (3.59) is much more compact and manifest with various analytic structures.

4. The general frame

Having the detail computations in the bubble, in this section we will set up the general frame to find generation functions for general one-loop integrals with $(n+1)$ propagators. The system has n external

momenta $K_i, i = 1, \dots, n$ and $(n+1)$ masses $M_j^2, j = 0, 1, \dots, n$. Using the auxiliary vector R we have $(n+1)$ **auxiliary scalar products (ASP)** $r = R \cdot R, p_i = K_i \cdot R, i = 1, \dots, n$. From the experience in the bubble, we know that these ASP's are not good variables to solve differential equations produced by $\partial_R \cdot \partial_R$ and $K_i \cdot \partial_R, i = 1, \dots, n$. Thus we will discuss how to find these good variables in the first subsection. Then we discuss the differential equations in these new variables in the second subsection and finally their solutions in the third subsection.

4.1 Finding good variables

We will denote the good variables by x and $y_i, i = 1, \dots, n$. To simplify the differential equations, we need to impose following conditions

$$(K_i \cdot \partial_R)x = 0, \quad (K_i \cdot \partial_R)y_j \sim \delta_{ij}, \quad \forall i, j = 1, \dots, n . \quad (4.1)$$

To see there is indeed a solution for (4.1), let us define the Gram matrix G and the row vector P^T as

$$G_{ij} = K_i \cdot K_j, \quad (P^T)_i = K_i \cdot R . \quad (4.2)$$

Putting $y_j = \sum_t \beta_{jt} p_t$ to (4.1), it is easy to see that the condition becomes

$$K_i \cdot \partial_R y_j = \sum_t \beta_{jt} (K_t \cdot K_i) \sim \delta_{ij} , \quad (4.3)$$

thus the matrix β can be solved as

$$\beta = |G|G^{-1} \quad (4.4)$$

where $|G|$ is the Gram determinant. For x , let us assume

$$x = |G|r + P^T A P, \quad A^T = A . \quad (4.5)$$

Since

$$K_i \cdot \partial_R x = |G|2p_i + 2(P^T)_{R \rightarrow K_i} A P, \quad (4.6)$$

where $P_{R \rightarrow K_i}$ means to replace vector R by the vector K_i . Collecting all i together, the right hand side of (4.6) is just $(2|G|I + 2GA)P$, thus we have the solution

$$A = -|G|G^{-1} . \quad (4.7)$$

Putting everything together, we finally have

$$x = |G|(r - P^T G^{-1} P), \quad Y = |G|G^{-1} P \quad (4.8)$$

where the mass dimensions of various quantities are

$$[|G|] = 2n, \quad [(G^{-1})_{ij}] = -2, \quad [A_{ij}] = 2(n-1), \quad [x] = 2(n+1), \quad [y_i] = 2n. \quad (4.9)$$

From (4.8) we can solve

$$P = \frac{1}{|G|} GY, \quad r = \frac{|G|x + Y^T GY}{|G|^2}. \quad (4.10)$$

4.2 The differential equations

Having found the good variables, we express differential operators $\partial_R \cdot \partial_R$ and $K_i \cdot \partial_R, i = 1, \dots, n$ using them. The first step is to use (4.8) to write

$$\frac{\partial}{\partial R^\mu} = (2|G|R_\mu - 2|G|\mathcal{K}_\mu^T G^{-1} P) \partial_x + |G|\mathcal{K}_\mu^T G^{-1} \partial_Y \quad (4.11)$$

where we have defined $\mathcal{K}^T = (K_1, \dots, K_n)$ and $\partial_Y^T = (\partial_{y_1}, \dots, \partial_{y_n})$. Thus we find

$$\mathcal{K} \cdot \frac{\partial}{\partial R^\mu} = |G|\partial_Y, \quad \partial_R \cdot \partial_R = 2|G|(D-n)\partial_x + 4|G|x\partial_x^2 + |G|^2 \partial_Y^T G^{-1} \partial_Y. \quad (4.12)$$

The differential equations for c_T , where T denotes different types of generation functions, have following pattern

$$K_i \cdot \partial_R c_T = \alpha_i c_T + H_{T;i}, \quad i = 1, 2, \dots, n \quad (4.13)$$

$$\partial_R \cdot \partial_R c_T = \alpha_R c_T + H_{T;R} \quad (4.14)$$

where α_R, α_i are constant (which are independent of T) and $H_{T;R}, H_{T;i}$ are known functions coming from lower topologies. Using the result (4.12) and (4.13), we find

$$|G|^2 \partial_Y^T G^{-1} \partial_Y c_T = \alpha_{\mathcal{K}}^T G^{-1} \alpha_{\mathcal{K}} c_T + H_{T;\mathcal{K}}^T G^{-1} \alpha_{\mathcal{K}} + |G|\partial_Y^T G^{-1} H_{T;\mathcal{K}} \quad (4.15)$$

where $\alpha_{\mathcal{K}}^T = (\alpha_1, \dots, \alpha_n)$ and $H_{T;\mathcal{K}}^T = (H_{T;1}, \dots, H_{T;n})$. Thus the differential equations (4.13) can be written as

$$\left(\partial_{y_i} - \frac{\alpha_i}{|G|} \right) c_T = \frac{1}{|G|} H_{T;i} \quad i = 1, 2, \dots, n, \quad (4.16)$$

while (4.14) becomes

$$(4|G|x\partial_x^2 + 2|G|(D-n)\partial_x + \tilde{\alpha}_R) c_T = \mathcal{H}_{T;R} \quad (4.17)$$

with

$$\tilde{\alpha}_R = \alpha_{\mathcal{K}}^T G^{-1} \alpha_{\mathcal{K}} - \alpha_R, \quad \mathcal{H}_{T;R} = -H_{T;\mathcal{K}}^T G^{-1} \alpha_{\mathcal{K}} - |G| \partial_Y^T G^{-1} H_{T;\mathcal{K}} + H_{T;R} . \quad (4.18)$$

Having given the differential equations (4.16) and (4.17), there is an important point to be mentioned. For (4.16) and (4.17) to have solution, functions H are not arbitrary, but must satisfy the **integrability conditions**, which are

$$\left(\partial_{y_j} - \frac{\alpha_j}{|G|} \right) H_{T;i} = \left(\partial_{y_i} - \frac{\alpha_i}{|G|} \right) H_{T;j}, \quad \forall i, j = 1, 2, \dots, n \quad (4.19)$$

and

$$(4|G|x\partial_x^2 + 2|G|(D-n)\partial_x + \tilde{\alpha}_R) \frac{1}{|G|} H_{T;i} = \left(\partial_{y_i} - \frac{\alpha_i}{|G|} \right) \mathcal{H}_{T;R}, \quad \forall i = 1, 2, \dots, n . \quad (4.20)$$

Differential equations (4.16) and (4.17) are the type of (A.1) and (A.14) respectively in the Appendix A, for which the solution has been presented. In next subsection, we will solve them analytically.

4.3 Analytic solution

In this part, we will present the necessary steps of solving above differential equations (4.16) and (4.17). Let us solve them one by one. For differential equation (4.16) with $i = 1$, using the result (A.5) in the Appendix A, we have

$$c_T(x, y) = e^{\frac{\alpha_1}{|G|} y_1} \left(F_T(x, y_2, \dots, y_n) + \frac{1}{|G|} \int_0^{y_1} dw_1 e^{-\frac{\alpha_1}{|G|} w_1} H_{T;1}(x, w_1, y_2, \dots, y_n) \right) \quad (4.21)$$

where $F_T(x, y_2, \dots, y_n)$ does not depend on the variable y_1 . Now we act with $\left(\partial_{y_2} - \frac{\alpha_2}{|G|} \right)$ at the both sides of (4.21) to get the differential equation

$$\begin{aligned} \left(\partial_{y_2} - \frac{\alpha_2}{|G|} \right) F_T(x, y_2, \dots, y_n) &= -\frac{1}{|G|} \int_0^{y_1} dw_1 e^{-\frac{\alpha_1}{|G|} w_1} \left(\partial_{y_2} - \frac{\alpha_2}{|G|} \right) H_{T;1}(x, w_1, y_2, \dots, y_n) \\ &\quad + e^{-\frac{\alpha_1}{|G|} y_1} \frac{1}{|G|} H_{T;2}(x, y_1, y_2, \dots, y_n) . \end{aligned} \quad (4.22)$$

Using (A.5) we find

$$\begin{aligned}
F_T(x, y_2, \dots, y_n) &= e^{\frac{\alpha_2}{|G|}y_2} F_T(x, y_3, \dots, y_n) + e^{\frac{\alpha_2}{|G|}y_2} e^{-\frac{\alpha_1}{|G|}y_1} \frac{1}{|G|} \int_0^{y_2} dw_2 e^{-\frac{\alpha_2}{|G|}w_2} H_{T;2}(x, y_1, w_2, \dots, y_n) \\
&- e^{\frac{\alpha_2}{|G|}y_2} \frac{1}{|G|} \int_0^{y_1} dw_1 e^{-\frac{\alpha_1}{|G|}w_1} \left(e^{-\frac{\alpha_2}{|G|}y_2} H_{T;1}(x, w_1, y_2, \dots, y_n) - H_{T;1}(x, w_1, y_2 = 0, \dots, y_n) \right). \quad (4.23)
\end{aligned}$$

Putting (4.23) back to (4.21) and doing some algebraic manipulations, we get

$$\begin{aligned}
c_T(x, y) &= e^{\frac{\alpha_1}{|G|}y_1} e^{\frac{\alpha_2}{|G|}y_2} F_T(x, y_3, \dots, y_n) \\
&+ e^{\frac{\alpha_2}{|G|}y_2} \frac{1}{|G|} \int_0^{y_2} dw_2 e^{-\frac{\alpha_2}{|G|}w_2} H_{T;2}(x, y_1, w_2, \dots, y_n) \\
&+ e^{\frac{\alpha_1}{|G|}y_1} e^{\frac{\alpha_2}{|G|}y_2} \frac{1}{|G|} \int_0^{y_1} dw_1 e^{-\frac{\alpha_1}{|G|}w_1} H_{T;1}(x, w_1, y_2 = 0, \dots, y_n) \quad (4.24)
\end{aligned}$$

Repeating above procedure with the action $(\partial_{y_2} - \frac{\alpha_2}{|G|})$ we can solve $F_T(x, y_3, \dots, y_n)$ and then find

$$\begin{aligned}
c_T(x, y) &= e^{\frac{\alpha_1}{|G|}y_1} e^{\frac{\alpha_2}{|G|}y_2} e^{\frac{\alpha_3}{|G|}y_3} F_T(x, y_4, \dots, y_n) \\
&+ e^{\frac{\alpha_3}{|G|}y_3} \frac{1}{|G|} \int_0^{y_3} dw_3 e^{-\frac{\alpha_3}{|G|}w_3} H_{T;3}(x, y_1, y_2, w_3, \dots, y_n) \\
&+ e^{\frac{\alpha_2}{|G|}y_2} e^{\frac{\alpha_3}{|G|}y_3} \frac{1}{|G|} \int_0^{y_2} dw_2 e^{-\frac{\alpha_2}{|G|}w_2} H_{T;2}(x, y_1, w_2, y_3 = 0, \dots, y_n) \\
&+ e^{\frac{\alpha_1}{|G|}y_1} e^{\frac{\alpha_2}{|G|}y_2} e^{\frac{\alpha_3}{|G|}y_3} \frac{1}{|G|} \int_0^{y_1} dw_1 e^{-\frac{\alpha_1}{|G|}w_1} H_{T;1}(x, w_1, y_2 = 0, y_3 = 0, \dots, y_n) \quad (4.25)
\end{aligned}$$

By checking (4.24) and (4.25) we can see that after solving n first order differential equations (4.16) we get

$$c_T(x, y_1, \dots, y_n) = e^{\frac{\sum_{i=1}^n \alpha_i y_i}{|G|}} F(x) + \mathcal{H}_{T;K} \quad (4.26)$$

with

$$\mathcal{H}_{T;K} = \frac{1}{|G|} \widetilde{\sum}_{i=n}^1 e^{\frac{\widetilde{\sum}_{j=n}^i \alpha_j y_j}{|G|}} \int_0^{y_i} dw_i e^{-\frac{\alpha_i w_i}{|G|}} H_{T;i}(x, y_1, \dots, y_{i-1}, w_i, 0, \dots, 0) \quad (4.27)$$

where for simplicity we have defined the sum $\widetilde{\sum}_{i=a}^b$ to mean to take the sum over $(a, a-1, a-2, \dots, b)$ with $a \geq b$.

Before going to solve the only unknown function $F(x)$, let us check that the form (4.26) does satisfy the differential equations (4.16). When acting with $(\partial_{y_k} - \frac{\alpha_k}{|G|})$ on the both sides, it is easy to see that the first term at the right hand side of (4.26) and the terms in $\mathcal{H}_{T;K}$ with $i < k$ give zero contributions since

they contain only the factor $e^{\frac{-\alpha_k y_k}{|G|}}$ depending on y_k . For the term $i = k$ in $\mathcal{H}_{T;K}$, the action gives

$$\frac{1}{|G|} e^{\frac{\sum_{j=n}^{k+1} \alpha_j y_j}{|G|}} H_{T;k}(x, y_1, \dots, y_{k-1}, y_k, 0, \dots, 0) . \quad (4.28)$$

For the term $i = k + 1$ in $\mathcal{H}_{T;K}$, the action gives

$$\begin{aligned} & \frac{1}{|G|} e^{\frac{\sum_{j=n}^{k+1} \alpha_j y_j}{|G|}} \int_0^{y_{k+1}} dw_{k+1} e^{\frac{-\alpha_{k+1} w_{k+1}}{|G|}} \left(\partial_{y_k} - \frac{\alpha_k}{|G|} \right) H_{T;k+1}(x, y_1, \dots, y_k, w_{k+1}, 0, \dots, 0) \\ &= \frac{1}{|G|} e^{\frac{\sum_{j=n}^{k+1} \alpha_j y_j}{|G|}} \int_0^{y_{k+1}} dw_{k+1} e^{\frac{-\alpha_{k+1} w_{k+1}}{|G|}} \left(\partial_{w_{k+1}} - \frac{\alpha_{k+1}}{|G|} \right) H_{T;k}(x, y_1, \dots, y_k, w_{k+1}, 0, \dots, 0) \end{aligned} \quad (4.29)$$

where in the second line we have used the integrability condition (4.19). After partial integration we get

$$-\frac{1}{|G|} e^{\frac{\sum_{j=n}^{k+1} \alpha_j y_j}{|G|}} H_{T;k}(x, y_1, \dots, y_{k-1}, y_k, 0, \dots, 0) + \frac{1}{|G|} e^{\frac{\sum_{j=n}^{k+2} \alpha_j y_j}{|G|}} H_{T;k}(x, y_1, \dots, y_k, y_{k+1}, 0, \dots, 0) . \quad (4.30)$$

The first term in (4.30) cancels the term in (4.28) and we are left with the second term in (4.30). Now the pattern is clear. The $i = k + 2$ term in $\mathcal{H}_{T;K}$ will produce two terms after using the integrability condition and partial integration, the first term will cancel the second term in (4.30), while the second term will be the form

$$\frac{1}{|G|} e^{\frac{\sum_{j=n}^{k+3} \alpha_j y_j}{|G|}} H_{T;k}(x, y_1, \dots, y_k, y_{k+1}, y_{k+2}, 0, \dots, 0) . \quad (4.31)$$

Continuing to the term $i = n$ in $\mathcal{H}_{T;K}$ we will be left with $\frac{1}{|G|} H_{T;k}(x, y_1, \dots, y_n)$, thus we have proved that (4.26) does satisfy the differential equations (4.16).

Now we consider the differential equation (4.17). Using the form (4.26), we derive

$$(4|G|x\partial_x^2 + 2|G|(D - n)\partial_x + \tilde{\alpha}_R) F(x) = e^{\frac{-\sum_{i=1}^n \alpha_i y_i}{|G|}} (\mathcal{H}_{T;R} - (4|G|x\partial_x^2 + 2|G|(D - n)\partial_x + \tilde{\alpha}_R) \mathcal{H}_{T;K}) \quad (4.32)$$

One important point of (4.32) is that the right hand side must be y_i -independent. To check this point, we act ∂_{y_k} at the right hand side to give

$$\begin{aligned} & -\frac{\alpha_k}{|G|} e^{\frac{-\sum_{i=1}^n \alpha_i y_i}{|G|}} (\mathcal{H}_{T;R} - (4|G|x\partial_x^2 + 2|G|(D - n)\partial_x + \tilde{\alpha}_R) \mathcal{H}_{T;K}) \\ & + e^{\frac{-\sum_{i=1}^n \alpha_i y_i}{|G|}} (\partial_{y_k} \mathcal{H}_{T;R} - (4|G|x\partial_x^2 + 2|G|(D - n)\partial_x + \tilde{\alpha}_R) \partial_{y_k} \mathcal{H}_{T;K}) . \end{aligned} \quad (4.33)$$

Since we have proved

$$\left(\partial_{y_k} - \frac{\alpha_k}{|G|}\right) \mathcal{H}_{T;K} = \frac{1}{|G|} H_{T;k} , \quad (4.34)$$

(4.33) is simplified to

$$-\frac{\alpha_k}{|G|} e^{-\frac{\sum_{i=1}^n \alpha_i y_i}{|G|}} \mathcal{H}_{T;R} + e^{-\frac{\sum_{i=1}^n \alpha_i y_i}{|G|}} \left(\partial_{y_k} \mathcal{H}_{T;R} - (4|G|x\partial_x^2 + 2|G|(D-n)\partial_x + \tilde{\alpha}_R) \frac{1}{|G|} H_{T;k} \right) . \quad (4.35)$$

Using the integrability condition (4.20) we get

$$-\frac{\alpha_k}{|G|} e^{-\frac{\sum_{i=1}^n \alpha_i y_i}{|G|}} \mathcal{H}_{T;R} + e^{-\frac{\sum_{i=1}^n \alpha_i y_i}{|G|}} \left(\partial_{y_k} \mathcal{H}_{T;R} - \left(\partial_{y_k} - \frac{\alpha_k}{|G|} \right) \mathcal{H}_{T;R} \right) = 0 . \quad (4.36)$$

Having checked the y -independent, we can take y_i to be any values at the right hand side of (4.32). From the expression (4.27) one can see that if we take $y_1 = y_2 = \dots = y_n = 0$, we have $\mathcal{H}_{T;K} = 0$, thus (4.32) is simplified to

$$(4|G|x\partial_x^2 + 2|G|(D-n)\partial_x + \tilde{\alpha}_R) F(x) = \mathcal{H}_{T;R}(x, 0, 0, \dots, 0) . \quad (4.37)$$

Above differential equation is the form of (A.14) in the Appendix A and we can write down the solution immediately (see (A.33))

$$F(x) = F_0(x) \left(f_0 + \int_0^x dw w^{-\frac{(D-n)}{2}} e^{-2F_0(w)} \int_0^w d\xi \frac{1}{A} \mathcal{H}_{T;R}(x, 0, 0, \dots, 0) F_0^{-1}(\xi) e^{2F_0(\xi)} \xi^{\frac{(D-n)}{2}-1} \right) \quad (4.38)$$

where

$$F_0(x) = {}_0F_1(\emptyset; \frac{(D-n)}{2}; \frac{-\tilde{\alpha}_R x}{4|G|}) . \quad (4.39)$$

Putting (4.39) back to (4.26) we get the final analytic expression of generation functions

$$c_T(x, y_1, \dots, y_n) = \mathcal{H}_{T;K} + e^{\frac{\sum_{i=1}^n \alpha_i y_i}{|G|}} F_0(x) \left(f_0 + \int_0^x dw w^{-\frac{(D-n)}{2}} e^{-2F_0(w)} \int_0^w d\xi \frac{1}{A} \mathcal{H}_{T;R}(x, 0, 0, \dots, 0) F_0(\xi)^{-1} e^{2F_0(\xi)} \xi^{\frac{(D-n)}{2}-1} \right) . \quad (4.40)$$

When we consider the generation functions of reduction coefficients of one-loop integrals with $(n+1)$ propagators, there is a special case, where all $H_{T;i}, H_{T;R}$ are zero. For this case, we can write down immediately the generation function

$$c_{n+1 \rightarrow n+1}(R, K_1, \dots, K_n) = {}_0F_1(\emptyset; \frac{(D-n)}{2}; \frac{-\tilde{\alpha}_R x}{4|G|}) e^{\frac{\sum_{i=1}^n \alpha_i y_i}{|G|}} . \quad (4.41)$$

5. Triangle

In this part we present another example, i.e., the triangle, to demonstrate the general frame laid down in previous section. The seven generation functions are defined by

$$\begin{aligned}
I_{tri}(t, R) &\equiv \int d\ell \frac{e^{t(2\ell \cdot R)}}{(\ell^2 - M_0^2)((\ell - K_1)^2 - M_1^2)((\ell - K_2)^2 - M_2^2)} \equiv \int d\ell \frac{e^{t(2\ell \cdot R)}}{D_0 D_1 D_2} \\
&= c_{3 \rightarrow 3} \int d\ell \frac{1}{D_0 D_1 D_2} + c_{3 \rightarrow 2; \hat{i}} \int d\ell \frac{1}{\prod_{j \neq i, 0}^2 D_j} + c_{3 \rightarrow 1; i} \int d\ell \frac{1}{D_i}.
\end{aligned} \tag{5.1}$$

Using the permutation symmetry and the shifting of loop momentum we can find nontrivial relations among these seven generation functions. The first group of relations is

$$\begin{aligned}
c_{3 \rightarrow 3}(t, R; M_0; K_1, M_1; K_2, M_2) &= c_{3 \rightarrow 3}(t, R; M_0; K_2, M_2; K_1, M_1), \\
c_{3 \rightarrow 2; \hat{0}}(t, R; M_0; K_1, M_1; K_2, M_2) &= c_{3 \rightarrow 2; \hat{0}}(t, R; M_0; K_2, M_2; K_1, M_1), \\
c_{3 \rightarrow 2; \hat{1}}(t, R; M_0; K_1, M_1; K_2, M_2) &= c_{3 \rightarrow 2; \hat{2}}(t, R; M_0; K_2, M_2; K_1, M_1), \\
c_{3 \rightarrow 2; \hat{2}}(t, R; M_0; K_1, M_1; K_2, M_2) &= c_{3 \rightarrow 2; \hat{1}}(t, R; M_0; K_2, M_2; K_1, M_1), \\
c_{3 \rightarrow 1; 0}(t, R; M_0; K_1, M_1; K_2, M_2) &= c_{3 \rightarrow 1; 0}(t, R; M_0; K_2, M_2; K_1, M_1), \\
c_{3 \rightarrow 1; 1}(t, R; M_0; K_1, M_1; K_2, M_2) &= c_{3 \rightarrow 1; 2}(t, R; M_0; K_2, M_2; K_1, M_1), \\
c_{3 \rightarrow 1; 2}(t, R; M_0; K_1, M_1; K_2, M_2) &= c_{3 \rightarrow 1; 1}(t, R; M_0; K_2, M_2; K_1, M_1).
\end{aligned} \tag{5.2}$$

The second group of relations is

$$\begin{aligned}
c_{3 \rightarrow 3}(t, R; M_0; K_1, M_1; K_2, M_2) &= e^{2K_1 \cdot R} c_{3 \rightarrow 3}(t, R; M_1; -K_1, M_0; K_2 - K_1, M_2), \\
c_{3 \rightarrow 2; \hat{0}}(t, R; M_0; K_1, M_1; K_2, M_2) &= e^{2K_1 \cdot R} c_{3 \rightarrow 2; \hat{1}}(t, R; M_1; -K_1, M_0; K_2 - K_1, M_2), \\
c_{3 \rightarrow 2; \hat{1}}(t, R; M_0; K_1, M_1; K_2, M_2) &= e^{2K_1 \cdot R} c_{3 \rightarrow 2; \hat{0}}(t, R; M_1; -K_1, M_0; K_2 - K_1, M_2), \\
c_{3 \rightarrow 2; \hat{2}}(t, R; M_0; K_1, M_1; K_2, M_2) &= e^{2K_1 \cdot R} c_{3 \rightarrow 2; \hat{1}}(t, R; M_1; -K_1, M_0; K_2 - K_1, M_2), \\
c_{3 \rightarrow 1; 0}(t, R; M_0; K_1, M_1; K_2, M_2) &= e^{2K_1 \cdot R} c_{3 \rightarrow 1; 1}(t, R; M_1; -K_1, M_0; K_2 - K_1, M_2), \\
c_{3 \rightarrow 1; 1}(t, R; M_0; K_1, M_1; K_2, M_2) &= e^{2K_1 \cdot R} c_{3 \rightarrow 1; 0}(t, R; M_1; -K_1, M_0; K_2 - K_1, M_2), \\
c_{3 \rightarrow 1; 2}(t, R; M_0; K_1, M_1; K_2, M_2) &= e^{2K_1 \cdot R} c_{3 \rightarrow 1; 1}(t, R; M_1; -K_1, M_0; K_2 - K_1, M_2).
\end{aligned} \tag{5.3}$$

The third group of relations is

$$\begin{aligned}
c_{3 \rightarrow 3}(t, R; M_0; K_1, M_1; K_2, M_2) &= e^{2K_2 \cdot R} c_{3 \rightarrow 3}(t, R; M_2; K_1 - K_2, M_1; -K_2, M_0), \\
c_{3 \rightarrow 2; \widehat{0}}(t, R; M_0; K_1, M_1; K_2, M_2) &= e^{2K_2 \cdot R} c_{3 \rightarrow 2; \widehat{2}}(t, R; M_2; K_1 - K_2, M_1; -K_2, M_0), \\
c_{3 \rightarrow 2; \widehat{1}}(t, R; M_0; K_1, M_1; K_2, M_2) &= e^{2K_2 \cdot R} c_{3 \rightarrow 2; \widehat{1}}(t, R; M_2; K_1 - K_2, M_1; -K_2, M_0), \\
c_{3 \rightarrow 2; \widehat{2}}(t, R; M_0; K_1, M_1; K_2, M_2) &= e^{2K_2 \cdot R} c_{3 \rightarrow 2; \widehat{0}}(t, R; M_2; K_1 - K_2, M_1; -K_2, M_0), \\
c_{3 \rightarrow 1; 0}(t, R; M_0; K_1, M_1; K_2, M_2) &= e^{2K_2 \cdot R} c_{3 \rightarrow 1; 2}(t, R; M_2; K_1 - K_2, M_1; -K_2, M_0), \\
c_{3 \rightarrow 1; 1}(t, R; M_0; K_1, M_1; K_2, M_2) &= e^{2K_2 \cdot R} c_{3 \rightarrow 1; 1}(t, R; M_2; K_1 - K_2, M_1; -K_2, M_0), \\
c_{3 \rightarrow 1; 2}(t, R; M_0; K_1, M_1; K_2, M_2) &= e^{2K_2 \cdot R} c_{3 \rightarrow 1; 0}(t, R; M_2; K_1 - K_2, M_1; -K_2, M_0).
\end{aligned} \tag{5.4}$$

Using these three groups of relations, we just need to compute three generation functions, for example, $c_{3 \rightarrow 3}$, $c_{3 \rightarrow 2; \widehat{1}}$ and $c_{3 \rightarrow 1; 0}$. The mass dimensions of them are

$$[t] = -2, \quad [c_{3 \rightarrow 3}] = 0, \quad [c_{3 \rightarrow 2; \widehat{i}}] = -2, \quad [c_{3 \rightarrow 1; i}] = -4. \tag{5.5}$$

5.1 The differential equations

Since we have given enough details in the section of bubble, here we will be more briefly. Using $\partial_R \cdot \partial_R, K_1 \cdot \partial_R, K_2 \cdot \partial_R$ operators, we can find

$$\partial_R \cdot \partial_R c_T(t, R; M_0; K_1, M_1; K_2, M_2) = 4t^2 M_0^2 c_T(t, R; M_0; K_1, M_1; K_2, M_2) + 4t^2 \xi_T h_T, \tag{5.6}$$

$$K_1 \cdot \partial_R c_T(t, R; M_0; K_1, M_1; K_2, M_2) = t f_1 c_T(t, R; M_0; K_1, M_1; K_2, M_2) - t \widetilde{\xi}_T \widetilde{h}_T + t \xi_T h_T, \tag{5.7}$$

$$K_2 \cdot \partial_R c_T(t, R; M_0; K_1, M_1; K_2, M_2) = t f_2 c_T(t, R; M_0; K_1, M_1; K_2, M_2) - t \widehat{\xi}_T \widehat{h}_T + t \xi_T h_T, \tag{5.8}$$

where T is the index for different generation functions and $f_1 = K_1^2 - M_1^2 + M_0^2$, $f_2 = K_2^2 - M_2^2 + M_0^2$. The various constant $\xi, \widetilde{\xi}, \widehat{\xi}$ are given in the table

T	ξ_T	h_T	$\widetilde{\xi}_T$	\widetilde{h}_T	$\widehat{\xi}_T$	\widehat{h}_T
$\{3 \rightarrow 3\}$	0	0	0	0	0	0
$\{3 \rightarrow 2; \widehat{0}\}$	1	$h_{3 \rightarrow 2; \widehat{0}}$	0	0	0	0
$\{3 \rightarrow 2; \widehat{1}\}$	0	0	1	$\widetilde{h}_{3 \rightarrow 2; \widehat{1}}$	0	0
$\{3 \rightarrow 2; \widehat{2}\}$	0	0	0	0	1	$\widehat{h}_{3 \rightarrow 2; \widehat{2}}$
$\{3 \rightarrow 1; 0\}$	0	0	1	$\widetilde{h}_{3 \rightarrow 1; 0}$	1	$\widehat{h}_{3 \rightarrow 1; 0}$
$\{3 \rightarrow 1; 1\}$	1	$h_{3 \rightarrow 1; 1}$	0	0	1	$\widehat{h}_{3 \rightarrow 1; 1}$
$\{3 \rightarrow 1; 2\}$	1	$h_{3 \rightarrow 1; 2}$	1	$\widetilde{h}_{3 \rightarrow 1; 2}$	0	0

(5.9)

while h_T are given by

$$\begin{aligned}
h_{3 \rightarrow 2; \hat{0}} &= e^{2K_1 \cdot R} c_{2 \rightarrow 2}(t, R^2, (K_2 - K_1) \cdot R, (K_2 - K_1)^2; M_1, M_2), \\
h_{3 \rightarrow 1; \hat{1}} &= e^{2K_1 \cdot R} c_{2 \rightarrow 2; \hat{1}}(t, R^2, (K_2 - K_1) \cdot R, (K_2 - K_1)^2; M_1, M_2), \\
h_{3 \rightarrow 1; \hat{2}} &= e^{2K_1 \cdot R} c_{2 \rightarrow 2; \hat{0}}(t, R^2, (K_2 - K_1) \cdot R, (K_2 - K_1)^2; M_1, M_2),
\end{aligned} \tag{5.10}$$

and \tilde{h}_T are given by

$$\begin{aligned}
\tilde{h}_{3 \rightarrow 2; \hat{1}} &= c_{2 \rightarrow 2}(t, R^2, K_2 \cdot R, K_2^2; M_0, M_2), \\
\tilde{h}_{3 \rightarrow 1; \hat{0}} &= c_{2 \rightarrow 1; \hat{1}}(t, R^2, K_2 \cdot R, K_2^2; M_0, M_2), \\
\tilde{h}_{3 \rightarrow 1; \hat{2}} &= c_{2 \rightarrow 1; \hat{0}}(t, R^2, K_2 \cdot R, K_2^2; M_0, M_2),
\end{aligned} \tag{5.11}$$

and \hat{h}_T are given by

$$\begin{aligned}
\hat{h}_{3 \rightarrow 2; \hat{2}} &= c_{2 \rightarrow 2}(t, R^2, K_1 \cdot R, K_1^2; M_0, M_1), \\
\hat{h}_{3 \rightarrow 1; \hat{0}} &= c_{2 \rightarrow 1; \hat{1}}(t, R^2, K_1 \cdot R, K_1^2; M_0, M_1), \\
\hat{h}_{3 \rightarrow 1; \hat{1}} &= c_{2 \rightarrow 1; \hat{0}}(t, R^2, K_1 \cdot R, K_1^2; M_0, M_1).
\end{aligned} \tag{5.12}$$

The differential equations (5.6), (5.7) and (5.8) are indeed the form (4.13) and (4.14) in previous section.

For triangle, the natural variables for generation functions are

$$r = R \cdot R, \quad p_1 = K_1 \cdot R, \quad p_2 = K_2 \cdot R. \tag{5.13}$$

However, the good variables for differential equations (5.6), (5.7) and (5.8) are x, y_1, y_2 as

$$\begin{aligned}
x &= [K_1^2 K_2^2 - (K_1 \cdot K_2)^2] r - K_2^2 p_1^2 - K_1^2 p_2^2 + (2K_1 \cdot K_2) p_1 p_2 \\
&= |G| r - p_1 y_1 - p_2 y_2, \\
y_2 &= -(K_1 \cdot K_2) p_1 + K_1^2 p_2, \quad y_1 = -(K_1 \cdot K_2) p_2 + K_2^2 p_1,
\end{aligned} \tag{5.14}$$

which are defined in (4.8) with Gram matrix

$$G(K_1, K_2) \equiv \begin{pmatrix} K_1^2 & K_1 \cdot K_2 \\ K_1 \cdot K_2 & K_2^2 \end{pmatrix}, \quad G^{-1} = \frac{1}{|G|} \begin{pmatrix} K_2^2 & -K_1 \cdot K_2 \\ -K_1 \cdot K_2 & K_1^2 \end{pmatrix}. \tag{5.15}$$

Using the new variables, the differential equations are

$$\left(\partial_{y_1} - \frac{\alpha_1}{|G|}\right) c_T = \frac{1}{|G|} H_{T;1}, \quad \left(\partial_{y_2} - \frac{\alpha_2}{|G|}\right) c_T = \frac{1}{|G|} H_{T;2} \quad (5.16)$$

$$(4|G|x\partial_x^2 + 2(D-2)|G|\partial_x + \tilde{\alpha}_R) c_T = \mathcal{H}_{T;R} \quad (5.17)$$

where

$$\alpha_1 = tf_1, \quad \alpha_2 = tf_2, \quad \tilde{\alpha}_R = \frac{t^2 f_1^2 K_2^2 + t^2 f_2^2 K_1^2 - 2t^2 f_1 f_2 (K_1 \cdot K_2)}{|G|} - 4t^2 M_0^2 \quad (5.18)$$

and

$$\begin{aligned} \mathcal{H}_{T;R} = & - \left(K_2^2 \partial_{y_1} - (K_1 \cdot K_2) \partial_{y_2} + \frac{\alpha_1 K_2^2 - \alpha_2 (K_1 \cdot K_2)}{|G|} \right) H_{T;1} \\ & - \left(K_1^2 \partial_{y_2} - (K_1 \cdot K_2) \partial_{y_1} + \frac{\alpha_2 K_1^2 - \alpha_1 (K_1 \cdot K_2)}{|G|} \right) H_{T;2} + H_{T;R}. \end{aligned} \quad (5.19)$$

The solution is given by

$$\begin{aligned} c_T(x, y_1, y_2) = & e^{\frac{\alpha_1}{|G|} y_1} e^{\frac{\alpha_2}{|G|} y_2} F_T^{tri}(x) + e^{\frac{\alpha_2}{|G|} y_2} \frac{1}{|G|} \int_0^{y_2} dw_2 e^{-\frac{\alpha_2}{|G|} w_2} H_{T;2}(x, y_1, w_2) \\ & + e^{\frac{\alpha_1}{|G|} y_1} e^{\frac{\alpha_2}{|G|} y_2} \frac{1}{|G|} \int_0^{y_1} dw_1 e^{-\frac{\alpha_1}{|G|} w_1} H_{T;1}(x, w_1, 0) \end{aligned} \quad (5.20)$$

where

$$F_0^{tri}(x) = {}_0F_1(\emptyset; \frac{(D-2)}{2}; \frac{-\tilde{\alpha}_R x}{4|G|}) \quad (5.21)$$

and

$$F^{tri}(x) = F_0^{tri}(x) \left(f_0 + \int_0^x dw w^{-\frac{(D-2)}{2}} e^{-2F_0^{tri}(w)} \int_0^w d\xi \frac{1}{A} \mathcal{H}_{T;R}(x, 0, 0, \dots, 0) F_0^{tri}(\xi)^{-1} e^{2F_0^{tri}(\xi)} \xi^{\frac{(D-2)}{2}-1} \right) \quad (5.22)$$

The generation function $c_{3 \rightarrow 3}$: For this one, from the Table (5.9), we see that $H_{T;1} = H_{T;2} = H_{T;R} = 0$, thus we write down immediately

$$c_{3 \rightarrow 3} = e^{\frac{\alpha_1}{|G|} y_1} e^{\frac{\alpha_2}{|G|} y_2} {}_0F_1(\emptyset; \frac{(D-2)}{2}; \frac{-\tilde{\alpha}_R x}{4|G|}) \quad (5.23)$$

The generation function $c_{3 \rightarrow 2}$: There are three generation functions $c_{3 \rightarrow 2; \hat{i}}$. We want to choose one of them with simplest H_1, H_2, H_R . Checking with Table (5.9), we see that if we consider $c_{3 \rightarrow 2; \hat{1}}$, we will

have $H_R = 0, H_2 = 0$ and

$$\begin{aligned} H_{3 \rightarrow 2; \hat{1}; 1}(x, y_1, y_2) &= -c_{2 \rightarrow 2}(t, R^2, K_2 \cdot R, K_2^2; M_0, M_2) \\ &= -{}_0F_1\left(\emptyset; \frac{D-1}{2}; \left(\frac{t^2(4K_2^2 M_0^2 - f_2^2)(K_2^2 r - p_2^2)}{4(K_2^2)^2}\right)\right) e^{\frac{t f_2}{K_2^2} p_2} \end{aligned} \quad (5.24)$$

where we have used the result (3.44) and expressed r, p_1, p_2 using (4.10)

$$p_1 = \frac{K_1^2 y_1 + (K_1 \cdot K_2) y_2}{|G|}, \quad p_2 = \frac{K_2^2 y_2 + (K_1 \cdot K_2) y_1}{|G|}, \quad r = \frac{|G|x + K_1^2 y_1^2 + K_2^2 y_2^2 + 2(K_1 \cdot K_2) y_1 y_2}{|G|^2}. \quad (5.25)$$

Thus by (5.19) we can find

$$\begin{aligned} \mathcal{H}_{3 \rightarrow 2; \hat{1}; R} &= \left(\frac{\alpha_1 K_2^2 - \alpha_2 (K_1 \cdot K_2)}{|G|}\right) e^{\frac{t f_2}{K_2^2} p_2} {}_0F_1\left(\emptyset; \frac{D-1}{2}; \left(\frac{t^2(4K_2^2 M_0^2 - f_2^2)(K_2^2 r - p_2^2)}{4(K_2^2)^2}\right)\right) \\ &+ \frac{t^2(4K_2^2 M_0^2 - f_2^2)}{4(K_2^2)^2} \frac{2K_2^2 y_1}{|G|} e^{\frac{t f_2}{K_2^2} p_2} {}_0F_1^{(1)}\left(\emptyset; \frac{D-1}{2}; \left(\frac{t^2(4K_2^2 M_0^2 - f_2^2)(K_2^2 r - p_2^2)}{4(K_2^2)^2}\right)\right) \end{aligned} \quad (5.26)$$

where we have defined

$${}_A F_B^{(n)}(a_1, \dots, a_A; b_1, \dots, b_B; x) = \frac{d^n}{dx^n} {}_A F_B(a_1, \dots, a_A; b_1, \dots, b_B; x) \quad (5.27)$$

Putting them back to (5.20) we can find the analytic expression. Using it we can get the explicit series expansion as discussed in the Appendix A.

For generation functions $c_{3 \rightarrow 1; i}$, we can do the similar calculations. The key is to find $H_{T; i}$ and $H_{T; R}$, which is the generation functions of one order lower topology. Thus we see the recursive structures of generation functions from lower topologies to higher topologies. The logic is clear although working out details takes some effects.

6. Conclusion

In this paper, we have introduced the concept of generation function for reduction coefficients of loop integrals. For one-loop integrals, using the recent proposal of auxiliary vector R , we can construct two types of differential operators $\frac{\partial}{\partial R} \cdot \frac{\partial}{\partial R}$ and $K_i \cdot \frac{\partial}{\partial R}$. Using these operators, we can establish corresponding differential equations for generation functions. By proper changing of variables, these differential equations can be written into the decoupled form, thus one can solve them one by one analytically. Obviously, one could try to apply the same idea to discuss the reduction problem for two and higher loop integrals. But with the appearance of irreducible scalar products, the problem become harder. One can try to use the IBP relation in the Baikov representation [43] as did in [40, 41].

Acknowledgments

We would like to thank Jiaqi Chen, Tingfei Li and Rui Yu for useful discussion and early participant. This work is supported by Chinese NSF funding under Grant No.11935013, No.11947301, No.12047502 (Peng Huanwu Center).

A. Solving differential equations

As shown in the paper, differential equations for generation functions can be reduced to two typical ones. In this appendix, we present details of solving these typical differential equations.

The first order differential equation: The first typical differential equation is following first order differential equation

$$\boxed{\left(A\frac{d}{dx} + B\right)F(x) = H(x)} \quad (\text{A.1})$$

where A, B are independent of x and $H(x)$ is the known function of x . To solve it, first we solve the homogenous part

$$\left(A\frac{d}{dx} + B\right)F_0(x) = 0, \quad \implies \quad F_0(x) = e^{-\frac{B}{A}x}. \quad (\text{A.2})$$

Then we write $F(x) = F_0(x)F_1(x)$ in (A.1) to get the differential equation for $F_1(x)$ as

$$A\frac{d}{dx}F_1(x) = F_0^{-1}H(x), \quad (\text{A.3})$$

thus we have

$$F_1(x) = F_1(x=0) + \int_0^x dt \frac{1}{A} e^{\frac{B}{A}t} H(t). \quad (\text{A.4})$$

Knowing the special solution of $F(x)$ in (A.4), the general solution of (A.1) is given by

$$\boxed{F(x) = F_0(x)F_1(x) + \tilde{\alpha}F_0(x) = e^{-\frac{B}{A}x} \left(\alpha + \int_0^x dw \frac{1}{A} e^{\frac{B}{A}w} H(w) \right)} \quad (\text{A.5})$$

where the constant α is determined by the boundary condition.

If we want to have the series expansion of x , there are several ways to do. The first one is to carry out

the integration

$$\int_0^x dt \frac{1}{A} e^{\frac{B}{A}t} H(t) = \frac{x}{A} \sum_{n=0}^{\infty} \sum_{a=0}^{\infty} h_n x^n \frac{1}{(n+a+1)} \frac{\left(\frac{Bx}{A}\right)^a}{a!} \quad (\text{A.6})$$

where we have used the expansion of $H(x) = \sum_{n=0}^{\infty} h_n x^n$. The second way is to put $F(x) = \sum_{n=0}^{\infty} f_n x^n$ directly to (A.1) to arrive the recursion relation

$$f_{n+1} = \frac{-B}{(n+1)A} f_n + \frac{h_n}{(n+1)A} \equiv \gamma(n) f_n + \rho(n), \quad n \geq 0 . \quad (\text{A.7})$$

The recursive relation (A.7) can be solved as

$$f_{n+1} = f_0 \prod_{i=0}^n \gamma(i) + \sum_{i=0}^n \rho(i) \prod_{j=i+1}^n \gamma(j) . \quad (\text{A.8})$$

It is easy to compute

$$\Xi[n+1] = \prod_{i=0}^n \gamma(i) = \frac{(-B)^{n+1}}{(n+1)! A^{n+1}} \quad (\text{A.9})$$

and

$$\rho(i) \prod_{j=i+1}^{n-1} \gamma(j) = \frac{h_i}{(i+1)A} \prod_{j=i+1}^{n-1} \gamma(j) = \frac{h_i}{-B} \prod_{j=i}^{n-1} \gamma(j) = \frac{h_i}{-B} \frac{\Xi[n]}{\Xi[i]} = \frac{h_i}{-B} \frac{i! (-B)^{n-i}}{n! A^{n-i}} , \quad (\text{A.10})$$

thus we have

$$F(x) = \sum_{n=0}^{\infty} \left(f_0 \frac{(-B)^n}{n! A^n} + \sum_{i=0}^{n-1} \frac{h_i i! (-B)^{n-1-i}}{n! A^{n-1-i}} \right) x^n . \quad (\text{A.11})$$

The third way is to use analytic expression (A.5). We need to compute $\frac{d^n F(x)}{dx^n}$ and then set $x = 0$. One can see that, for example,

$$\frac{dF(x)}{dx} = -\frac{B}{A} e^{-\frac{B}{A}x} \left(\alpha + \int_0^x dw \frac{1}{A} e^{\frac{B}{A}w} H(w) \right) + \frac{1}{A} H(x) , \quad (\text{A.12})$$

thus

$$\left. \frac{dF(x)}{dx} \right|_{x=0} = -\frac{B}{A} \alpha + \frac{1}{A} H(x=0) . \quad (\text{A.13})$$

The important point is that when $x = 0$ the integration $\int_0^x dw \dots = 0$, thus we have got rid of integration and all we need to do are the differentiations over $e^{\frac{B}{A}x}$ and $H(x)$.

The second order differential equation: The second typical differential equation met in this paper is the following second order differential equation

$$\boxed{\left(Ax \frac{d^2}{dx^2} + B \frac{d}{dx} + C \right) F(x) = H(x)} \quad (\text{A.14})$$

where A, B, C are independent of x and $H(x)$ is a known function of x . Let us to solve it using the series expansion. Writing

$$F(x) = \sum_{n=0}^{\infty} f_n x^n, \quad H(x) = \sum_{n=0}^{\infty} h_n x^n \quad (\text{A.15})$$

and putting back to (A.14) we have

$$\sum_{n=0}^{\infty} h_n x^n = \sum_{n=0}^{\infty} (An(n+1)f_{n+1}x^n + B(n+1)f_{n+1}x^n + Cf_n x^n), \quad (\text{A.16})$$

thus we have the relation

$$h_n = (n+1)(B+An)f_{n+1} + Cf_n, \quad n \geq 0. \quad (\text{A.17})$$

Using it, we can solve⁶

$$f_{n+1} = \frac{h_n}{(n+1)(B+An)} - C \frac{1}{(n+1)(B+An)} f_n \equiv \gamma(n)f_n + \rho(n), \quad n \geq 0. \quad (\text{A.18})$$

The recursive relation (A.18) can be solved as

$$f_{n+1} = f_0 \prod_{i=0}^n \gamma(i) + \sum_{i=0}^n \rho(i) \prod_{j=i+1}^n \gamma(j). \quad (\text{A.19})$$

It is easy to compute

$$\Xi[n+1] = \prod_{i=0}^n \gamma(i) = \frac{(-C)^{n+1}}{(n+1)!A^{n+1} \prod_{i=0}^n (B/A+i)} = \frac{(-C)^{n+1}}{(n+1)!A^{n+1} \left(\frac{B}{A}\right)_{n+1}} \quad (\text{A.20})$$

⁶An important point is that although (A.14) is second order differential equations, but because the x in front of $\frac{d^2}{dx^2}$, around $x = 0$ it is essentially the first order differential equation. This explains why using only f_0 and known $H(x)$ we can determine $F(x)$ using (A.18).

where we have used the **Pochhammer symbol** to simplify the expression⁷

$$(x)_n = \frac{\Gamma(x+n)}{\Gamma(x)} = \prod_{i=1}^n (x+(i-1)) . \quad (\text{A.21})$$

Using

$$\rho(i) \prod_{j=i+1}^{n-1} \gamma(j) = \frac{h_i}{(i+1)(B+Ai)} \prod_{j=i+1}^{n-1} \gamma(j) = \frac{-h_i}{C} \prod_{j=i}^{n-1} \gamma(j) = \frac{-h_i}{C} \frac{\Xi[n]}{\Xi[i]} \quad (\text{A.22})$$

we have

$$F(x) = \sum_{n=0}^{\infty} f_n x^n = \sum_{n=0}^{\infty} x^n \Xi[n] \left\{ f_0 + \sum_{i=0}^{n-1} \frac{-h_i}{C \Xi[i]} \right\} . \quad (\text{A.23})$$

Let us define the following function

$$H_{A,B,C}(x) = \sum_{i=0}^{\infty} \frac{-h_i x^i}{C \Xi[i]} = \sum_{i=0}^{\infty} \frac{i! A^i}{(-C)^{i+1}} \left(\frac{B}{A} \right)_i h_i x^i \quad (\text{A.24})$$

which can be considered as the "dual function" of $H(x)$ corresponding to the differential equation (A.14), then we can write

$$\sum_{i=0}^{n-1} \frac{-h_i}{C \Xi[i]} = [H_{A,B,C}(x=1)]_{x^{n-1}} \quad (\text{A.25})$$

where the symbol $[Y(x)]_{x^{n-1}}$ means to keep the Taylor series up to the order of $x^{(n-1)}$. Thus $F(x)$ can be written compactly as

$$F(x) = \sum_{n=0}^{\infty} f_n x^n = \sum_{n=0}^{\infty} x^n \Xi[n] \{ f_0 + [H_{A,B,C}(x=1)]_{x^{n-1}} \} . \quad (\text{A.26})$$

For the special case $H(x) = 0$, it is easy to see that

$$F(x) = f_0 F_0, \quad F_0(x) \equiv \sum_{n=0}^{\infty} \frac{(-C)^n x^n}{n! A^n \left(\frac{B}{A} \right)_n} . \quad (\text{A.27})$$

The expression (A.27) is nothing, but the special case of **generalized hypergeometric function** (see

⁷From the definition one can see that $(x)_{n=0} = 1, \forall x$.

(C.2) of [11]), which is defined as

$${}_A F_B(a_1, \dots, a_A; b_1, \dots, b_B; x) = \sum_{n=0}^{\infty} \frac{(a_1)_n \dots (a_A)_n x^n}{(b_1)_n \dots (b_B)_n n!}, \quad (\text{A.28})$$

thus we have

$$F(x) = f_0 F_0(x), \quad F_0 = {}_0 F_1\left(\emptyset; \frac{B}{A}; \frac{-Cx}{A}\right). \quad (\text{A.29})$$

Solution in (A.23) is given in the series expansion. We can also write it in the analytic expression. Writing $F(x) = F_0(x)F_1(x)$ we can find the differential equation of $F_1(x)$ as

$$H(x) = F_0(x) \left(Ax \frac{d^2}{dx^2} + (B + 2Ax \frac{dF_0(x)}{dx}) \frac{d}{dx} \right) F_1(x), \quad (\text{A.30})$$

which is the first order differential equation of $\frac{dF_1(x)}{dx} = U(x)$. Using the similar method as for the differential equation (A.1) we can solve

$$U(x) = x^{-\frac{B}{A}} e^{-2F_0(x)} \left(\alpha_1 + \int_0^x dw \frac{1}{A} H(w) F_0^{-1}(w) e^{2F_0(w)} w^{\frac{B}{A}-1} \right), \quad (\text{A.31})$$

thus we obtain

$$F(x) = F_0(x) \left(\alpha_2 + \int_0^x dw w^{-\frac{B}{A}} e^{-2F_0(w)} \left(\alpha_1 + \int_0^w d\xi \frac{1}{A} H(\xi) F_0^{-1}(\xi) e^{2F_0(\xi)} \xi^{\frac{B}{A}-1} \right) \right) \quad (\text{A.32})$$

where the α_1, α_2 can be determined using the initial condition of $F(x=0)$ and $\frac{dF(x=0)}{dx}$. Using the expansion of $F(x)$ we see that $\alpha_2 = f_0$ and $\alpha_1 = 0$ thus we have

$$\boxed{F(x) = F_0(x) \left(f_0 + \int_0^x dw w^{-\frac{B}{A}} e^{-2F_0(w)} \int_0^w d\xi \frac{1}{A} H(\xi) F_0^{-1}(\xi) e^{2F_0(\xi)} \xi^{\frac{B}{A}-1} \right)}. \quad (\text{A.33})$$

The analytic form (A.33) is very compact, but it is hard to carry out the integration in general. However, we can use it to get series expansion just like before. One can see that

$$\begin{aligned} \frac{dF(x)}{dx} &= \frac{dF_0(x)}{dx} \left(f_0 + \int_0^x dw w^{-\frac{B}{A}} e^{-2F_0(w)} \int_0^w dt \frac{1}{A} H(t) F_0^{-1}(t) e^{2F_0(t)} t^{\frac{B}{A}-1} \right) \\ &\quad + F_0(x) x^{-\frac{B}{A}} e^{-2F_0(x)} \int_0^x dt \frac{1}{A} H(t) F_0^{-1}(t) e^{2F_0(t)} t^{\frac{B}{A}-1} \end{aligned} \quad (\text{A.34})$$

When taking the value at the $x = 0$, we need to be careful for the second line and evaluate it as

$$\begin{aligned} & \int_0^x dt \frac{1}{A} H(t) F_0(t)^{-1} e^{2F_0(t)} t^{\frac{B}{A}-1} \rightarrow \frac{1}{A} H(x=0) F_0^{-1}(x=0) e^{2F_0(x=0)} \int_0^x dt t^{\frac{B}{A}-1} \\ & = \frac{1}{A} H(x=0) F_0^{-1}(x=0) e^{2F_0(x=0)} \frac{A}{B} x^{\frac{B}{A}} \Big|_0^x = H(x=0) F_0^{-1}(x=0) e^{2F_0(x=0)} \frac{1}{B} x^{\frac{B}{A}} \end{aligned} \quad (\text{A.35})$$

With the result (A.35), (A.34) gives

$$\frac{dF(x)}{dx} = f_0 \frac{dF_0(x)}{dx} \Big|_{x=0} + \frac{1}{B} H(x=0) \quad (\text{A.36})$$

From (A.14) we have

$$\frac{dF}{dx} \Big|_{x=0} = \frac{1}{B} H(x=0) - \frac{C}{B} F(x=0) \quad (\text{A.37})$$

thus we see that

$$f_0 = -\frac{C}{B} F(x=0) \left(\frac{dF_0(x)}{dx} \Big|_{x=0} \right)^{-1} \quad (\text{A.38})$$

For general $\frac{d^n F(x)}{dx^n}$ we do same thing to get the series expansion of x . At each step, we use (A.35) and there is no integration to be done for $x = 0$.

B. The explicit solutions of $c_{n,m}$ for bubble reduction

In this part, we will show how to get explicit solutions for the recursion relations (3.30), (3.31), (3.32) and (3.33).

B.1 The generation function $c_{2 \rightarrow 2}$

For this case, we have $h_T = 0$, i.e., all $h_{n,m} = 0$ in (3.30), (3.31), (3.32) and (3.33). Using (3.30) it is easy to find

$$c_{N,0} = \prod_{n=1}^N \frac{(-f^2 + 4K^2 M_0^2) t^2}{2K^2 n(D + 2n - 3)} = \prod_{n=0}^{N-1} \frac{K^2(\beta - \alpha^2)}{2(n+1)(D + 2n - 1)} \quad (\text{B.1})$$

where we have used the initial condition $c_{0,0} = 1$ and defined

$$\alpha = \frac{tf}{K^2}, \quad \beta = \frac{4t^2 M_0^2}{K^2}. \quad (\text{B.2})$$

Using (3.32) to compute first few $c_{n,m}$, one can see the pattern

$$c_{N,m} = \frac{1}{m!} d_{N,m} c_{N,0} \quad (\text{B.3})$$

where $d_{N,m}$ depends on both N, m and first few $d_{N,m}$ are

$$d_{N,0} = 1, \quad d_{N,1} = \alpha, \quad d_{N,2} = \frac{(D+2N)\alpha^2 - \beta}{(D+2N-1)}. \quad (\text{B.4})$$

Putting the form (B.3) to (3.32) we get the recursion relation

$$d_{N,m+1} = \frac{\beta}{\alpha} d_{N,m} - \frac{(\beta - \alpha^2)(D+2N+m-1)}{\alpha(D+2N-1)} d_{N+1,m} \quad (\text{B.5})$$

Using the formally defined operator \widehat{P} such that

$$\widehat{P}f(N, m) = f(N+1, m) \quad (\text{B.6})$$

the solution of (B.5) can be formally given by

$$\begin{aligned} d_{N,M} &= \prod_{m=1}^M \left(\frac{\beta}{\alpha} - \frac{(\beta - \alpha^2)(D+2N+m-2)}{\alpha(D+2N-1)} \widehat{P} \right), \quad M \geq 1, \\ &:= \left(\frac{\beta}{\alpha} - \frac{(\beta - \alpha^2)(D+2N+M-2)}{\alpha(D+2N-1)} \widehat{P} \right) \left(\frac{\beta}{\alpha} - \frac{(\beta - \alpha^2)(D+2N+(M-1)-2)}{\alpha(D+2N-1)} \widehat{P} \right) \\ &\quad \times \dots \left(\frac{\beta}{\alpha} - \frac{(\beta - \alpha^2)(D+2N+2-2)}{\alpha(D+2N-1)} \widehat{P} \right) \left(\frac{\beta}{\alpha} - \frac{(\beta - \alpha^2)(D+2N+1-2)}{\alpha(D+2N-1)} \widehat{P} \right) d_{N,0} \end{aligned} \quad (\text{B.7})$$

where since the appearance of the operator \widehat{P} , the ordering of the multiple factors is given explicitly. With a little computation, one can see that

$$d_{N,M} = \sum_{i=0}^M \frac{(-)^i \beta^{M-i} (\beta - \alpha^2)^i}{\alpha^M} d_{N,M;i} \quad (\text{B.8})$$

where

$$d_{N,M;i} = \sum_{1;[t_1, \dots, t_i]}^M \prod_{s=1}^i \frac{D+2(N+s-1) + (M+1-t_s) - 2}{D+2(N+s-1) - 1}, \quad i \geq 1; \quad d_{N,M;0} = 1 \quad (\text{B.9})$$

and the summation sign is defined as

$$\sum_{1 \leq t_1 < t_2 < \dots < t_i \leq M} := \sum_{1;[t_1, \dots, t_i]}^M \quad (\text{B.10})$$

Combining (B.3) with (B.1) and (B.8), we have the explicit solution for the generation function of $c_{2 \rightarrow 2}(t, r, p, K^2; M_0, M_1)$.

B.1.1 The generation function $c_{2 \rightarrow 1; \hat{1}}$

For this case, we have $h_T = c_{1 \rightarrow 1}(t, r, M_0)$. Using (2.5) to (3.30) one can write down

$$c_{n+1,0} = \gamma(n)c_{n,0} + \rho(n) \quad (\text{B.11})$$

with

$$\gamma(n) = \frac{K^2(\beta - \alpha^2)}{2(n+1)(D+2n-1)}, \quad \rho(n) = \frac{t^{2n+1}(M_0^2)^n \alpha}{2(D+2n-1)(n+1)! \left(\frac{D}{2}\right)_n} \quad (\text{B.12})$$

thus we can solve

$$c_{n+1,0} = c_{0,0} \prod_{i=0}^n \gamma(i) + \sum_{i=0}^n \rho(i) \prod_{j=i+1}^n \gamma(j) \quad (\text{B.13})$$

where for current case, the initial condition is $c_{0,0} = 0$. To find $c_{n,m}$ we write

$$c_{N,m} = \frac{1}{m!} d_{N,m} c_{N,0} - \frac{1}{m!} \frac{t \beta^N (K^2)^{N-1}}{N! 4^N \left(\frac{D}{2}\right)_N} b_{N,m} \quad (\text{B.14})$$

with the first few $d_{N,m}$ and $b_{N,m}$

$$\begin{aligned} d_{N,0} &= 1, & b_{N,0} &= 0 \\ d_{N,1} &= \alpha, & b_{N,1} &= 1 \\ d_{N,2} &= \frac{-\beta + \alpha^2(D+2N)}{(D+2N-1)}, & b_{N,2} &= \frac{\alpha(D+2N)}{(D+2N-1)} \end{aligned} \quad (\text{B.15})$$

Using the form (B.14) to (3.32) we get the recursion relations

$$d_{N,m+1} = \frac{\beta}{\alpha} d_{N,m} - \frac{(D+2N+m-1)(\beta - \alpha^2)}{\alpha(D+2N-1)} d_{N+1,m} \quad (\text{B.16})$$

$$b_{N,m+1} = \frac{\beta}{\alpha} b_{N,m} - \frac{\beta(D+2N+m-1)}{\alpha(D+2N)} b_{N+1,m} + \frac{(D+2N+m-1)}{(D+2N-1)} d_{N+1,m} \quad (\text{B.17})$$

The solution of $d_{N,m}$ can be similarly solved using the operator language as in previous subsection and we find

$$d_{N,M} = \sum_{i=0}^M \frac{(-)^i \beta^{M-i} (\beta - \alpha^2)^i}{\alpha^M} d_{N,M;i} \quad (\text{B.18})$$

with

$$d_{N,M;i} = \sum_{1:[t_1,\dots,t_i]}^M \prod_{s=1}^i \frac{D + 2(N + s - 1) + (M + 1 - t_s) - 2}{D + 2(N + s - 1) - 1}, \quad i \geq 1; \quad d_{N,M;0} = 1 \quad (\text{B.19})$$

The solution for $b_{N,m}$ is a little bit complicated because the third term at the right hand side of (B.17). To solve it, we write

$$b_{N,m+1} = \tilde{\gamma}_N(m)b_{N,m} + \tilde{\rho}_N(m) \quad (\text{B.20})$$

where

$$\tilde{\gamma}_N(m) = \left(\frac{\beta}{\alpha} - \frac{\beta(D + 2N + m - 1)}{\alpha(D + 2N)} \hat{P} \right), \quad \tilde{\rho}_N(m) = \frac{(D + 2N + m - 1)}{(D + 2N - 1)} d_{N+1,m} \quad (\text{B.21})$$

Iterating (B.20) with proper ordering we have

$$\begin{aligned} b_{N,m+1} &= \tilde{\gamma}_N(m)\tilde{\gamma}_N(m-1)\dots\tilde{\gamma}_N(0)b_{N,0} + \tilde{\gamma}_N(m)\tilde{\gamma}_N(m-1)\dots\tilde{\gamma}_N(1)\tilde{\rho}_N(0) \\ &\quad + \tilde{\gamma}_N(m)\tilde{\gamma}_N(m-1)\dots\tilde{\gamma}_N(2)\tilde{\rho}_N(1) + \dots + \tilde{\gamma}_N(m)\tilde{\rho}_N(m-1) + \tilde{\rho}_N(m) \end{aligned} \quad (\text{B.22})$$

Using

$$(\hat{P})^a \tilde{\rho}_N(m) = \frac{(D + 2(N + a) + m - 1)}{(D + 2(N + a) - 1)} d_{N+1+a,m} \quad (\text{B.23})$$

and

$$\begin{aligned} &\tilde{\gamma}_N(m)\tilde{\gamma}_N(m-1)\dots\tilde{\gamma}_N(m-k)\tilde{\rho}_N(m-k-1) \\ &= \sum_{i=0}^{k+1} (-)^i \left(\frac{\beta}{\alpha} \right)^{k+1} \left(\sum_{1:[t_1,\dots,t_i]}^{k+1} \prod_{s=1}^i \frac{D + 2(N + s - 1) + (m + 1 - t_s) - 1}{D + 2(N + s - 1)} \right) \\ &\quad \times \frac{(D + 2(N + i) + (m - k - 1) - 1)}{(D + 2(N + i) - 1)} d_{N+1+i,m-k-1} \end{aligned} \quad (\text{B.24})$$

we finally reach

$$\begin{aligned}
b_{N,m+1} &= \frac{(D+2N+m-1)}{(D+2N-1)} d_{N+1,m} + \sum_{k=0}^{m-1} \left(\frac{\beta}{\alpha}\right)^{k+1} \sum_{i=0}^{k+1} (-)^i \\
&\times \frac{(D+2(N+i)+(m-k-1)-1)}{(D+2(N+i)-1)} d_{N+1+i,m-k-1} \\
&\times \left(\sum_{1;[t_1,\dots,t_i]}^{k+1} \prod_{s=1}^i \frac{D+2(N+s-1)+(m+1-t_s)-1}{D+2(N+s-1)} \right) \tag{B.25}
\end{aligned}$$

where the condition $b_{N,0} = 0$ has been used. The formula (B.14) plus (B.13), (B.18) and (B.25) gives the explicit solution for generation function $c_{2 \rightarrow 1; \hat{1}}$.

Knowing $c_{2 \rightarrow 1; \hat{1}}$, we can use (3.5) to get the generation function $c_{2 \rightarrow 1; \hat{0}}$ or directly compute it using (3.30), (3.31), (3.32) and (3.33).

B.1.2 The proof of one useful relation

When we use the improved PV-reduction method with auxiliary vector R to discuss the reduction of sunset topology, an important reduction relation between different tensor ranks has been observed in [38]. Later this relation has been studied in [39, 40, 42]. For bubble it is given explicitly by

$$\begin{aligned}
I_2^{(r)} &= \frac{(D+2r-4)fp}{(D+r-3)K^2} I_2^{(r-1)} - \frac{(r-1)(4M_0^2 p^2 + (f^2 - 4M_0^2 K^2)r)}{(D+r-3)K^2} I_2^{(r-2)} \\
&+ \frac{p}{K^2} I_{2;\hat{0}}^{(r-1)} + \frac{(r-1)((K^2 - M_0^2 + M_1^2)r - 2p^2)}{(D+r-3)K^2} I_{2;\hat{0}}^{(r-2)} + \frac{-p}{K^2} I_{2;\hat{1}}^{(r-1)} + \frac{(r-1)fr}{(D+r-3)K^2} I_{2;\hat{1}}^{(r-2)} \tag{B.26}
\end{aligned}$$

where

$$I_2^{(r)} = \int d\ell \frac{(2\ell \cdot R)^r}{(\ell^2 - M_0^2)((\ell - K)^2 - M_1^2)}, \quad I_{2;\hat{1}}^{(r)} = \int d\ell \frac{(2\ell \cdot R)^r}{(\ell^2 - M_0^2)}, \quad I_{2;\hat{0}}^{(r)} = \int d\ell \frac{(2\ell \cdot R)^r}{(\ell - K)^2 - M_1^2} \tag{B.27}$$

We can use the series form to check the relation (B.26). Let us define

$$F[N+2] = c^{(N+2)} - \left(\frac{(D+2N)fp c^{(N+1)} - (N+1)(4M_0^2 p^2 + (f^2 - 4M_0^2 K^2)r) c^{(N)}}{(D+N-1)K^2} \right) \tag{B.28}$$

where

$$c^{(N)} = \frac{N!}{t^N} \sum_{n=0}^{[N/2]} c_{n,N-2n} r^n p^{N-2n} \tag{B.29}$$

and c can be $c_{2 \rightarrow 2}$ or $c_{2 \rightarrow 1}$. Depending on if N is even or odd, the computation details have some differences,

thus we consider N even only, while N odd will be similar. Expanding (B.28) we have

$$\begin{aligned}
F[2N+2] = & \frac{(2N+2)!}{t^{2N+2}} r^0 p^{2N+2} \left(c_{0,2N+2} - \left(\frac{(D+4N)tf c_{0,2N+1}}{(2N+2)(D+2N-1)K^2} - \frac{4M_0^2 t^2 c_{0,2N}}{2(N+1)(D+2N-1)K^2} \right) \right) \\
& + \frac{(2N+2)!}{t^{2N+2}} r^{N+1} p^0 \left(c_{N+1,0} - \left(-\frac{t^2(f^2 - 4M_0^2 K^2)}{(2N+2)(D+2N-1)K^2} c_{N,0} \right) \right) \\
& + \frac{(2N+2)!}{t^{2N+2}} \sum_{n=1}^N r^n p^{2N+2-2n} \left(c_{n,2N+2-2n} - \left(\frac{(D+4N)tf}{(2N+2)(D+2N-1)K^2} c_{n,2N+1-2n} \right. \right. \\
& \left. \left. - \frac{4M_0^2 t^2}{(2N+2)(D+2N-1)K^2} c_{n,2N-2n} - \frac{t^2(f^2 - 4M_0^2 K^2)}{(2N+2)(D+2N-1)K^2} c_{n-1,2N+2-2n} \right) \right) \quad (B.30)
\end{aligned}$$

For the term with $r^0 p^{2N+2}$, using the relation (3.29) with $n=0, m=2N$, we find the part inside the bracket is simplified to

$$\frac{-4\xi_R t^2 h_{0,2N} + \xi_K t(D+4N)h_{0,2N+1}}{2(N+1)(D+2N-1)K^2} \quad (B.31)$$

For the term with $r^{N+1} p^0$, using the relation (3.30) with $n=N$, we find the part inside the bracket is simplified to

$$-\frac{t^2(\xi_K f - \xi_R 4K^2)h_{N,0} + \xi_K t K^2 h_{N,1}}{2(N+1)(D+2N-1)K^2} \quad (B.32)$$

For the term with $r^n p^{2N+2-2n}$, the computation is a little bit complicated. First we use (3.28) with $n \rightarrow n-1$ and $m \rightarrow 2N-2n+2, 2N-2n+1, 2N-2n$ to write all $c_{i,j}$ with $i=n-1$. Then we use (3.29) with $n \rightarrow n-1$ and $m \rightarrow 2N-2n+1, 2N-2n$. After doing above two steps and making algebraic simplification, we get

$$\begin{aligned}
& \frac{t^3(\xi_K M_0^2 - \xi_R f)}{(N+1)(D+2N-1)nK^2} h_{n-1,2N-2n+1} + \frac{t^2(-\xi_K n f + \xi_R 4K^2(N+1))}{2(N+1)(D+2N-1)nK^2} h_{n-1,2N-2n+2} \\
& + \frac{-\xi_K(2N-2n+3)t}{2n(D+2N-1)} h_{n-1,2N-2n+3} \quad (B.33)
\end{aligned}$$

For different c we use the different known h_T as given in (3.17), (3.18) and (3.19), thus one can see the relation (B.26) is satisfied.

References

- [1] Z. Bern *et al.* [NLO Multileg Working Group], ‘‘The NLO multileg working group: Summary report,’’ [arXiv:0803.0494 [hep-ph]].

- [2] Johannes M. Henn and Jan C. Plefka, “Scattering Amplitudes in Gauge Theories”. Lecture Notes in Physics, Volume 883, Springer.
- [3] Henriette Elvang and Yu-Tin Huang, “Scattering Amplitudes in Gauge Theory and Gravity”. Cambridge University Press.
- [4] Z. Bern, L. J. Dixon, D. C. Dunbar and D. A. Kosower, “One loop n point gauge theory amplitudes, unitarity and collinear limits,” Nucl. Phys. B **425**, 217-260 (1994) doi:10.1016/0550-3213(94)90179-1 [arXiv:hep-ph/9403226 [hep-ph]].
- [5] Z. Bern, L. J. Dixon, D. C. Dunbar and D. A. Kosower, “Fusing gauge theory tree amplitudes into loop amplitudes,” Nucl. Phys. B **435**, 59-101 (1995) doi:10.1016/0550-3213(94)00488-Z [arXiv:hep-ph/9409265 [hep-ph]].
- [6] F. Cachazo, P. Svrcek and E. Witten, “MHV vertices and tree amplitudes in gauge theory,” JHEP **09**, 006 (2004) doi:10.1088/1126-6708/2004/09/006 [arXiv:hep-th/0403047 [hep-th]].
- [7] R. Britto, F. Cachazo and B. Feng, “Generalized unitarity and one-loop amplitudes in N=4 super-Yang-Mills,” Nucl. Phys. B **725**, 275-305 (2005) doi:10.1016/j.nuclphysb.2005.07.014 [arXiv:hep-th/0412103 [hep-th]].
- [8] R. Britto, F. Cachazo and B. Feng, “New recursion relations for tree amplitudes of gluons,” Nucl. Phys. B **715**, 499-522 (2005) doi:10.1016/j.nuclphysb.2005.02.030 [arXiv:hep-th/0412308 [hep-th]].
- [9] R. Britto, F. Cachazo, B. Feng and E. Witten, “Direct proof of tree-level recursion relation in Yang-Mills theory,” Phys. Rev. Lett. **94**, 181602 (2005) doi:10.1103/PhysRevLett.94.181602 [arXiv:hep-th/0501052 [hep-th]].
- [10] R. K. Ellis, Z. Kunszt, K. Melnikov and G. Zanderighi, “One-loop calculations in quantum field theory: from Feynman diagrams to unitarity cuts,” Phys. Rept. **518**, 141-250 (2012) doi:10.1016/j.physrep.2012.01.008 [arXiv:1105.4319 [hep-ph]].
- [11] S. Weinzierl, “Feynman Integrals,” [arXiv:2201.03593 [hep-th]].
- [12] G. Ossola, C. G. Papadopoulos and R. Pittau, “Reducing full one-loop amplitudes to scalar integrals at the integrand level,” Nucl. Phys. B **763**, 147-169 (2007) doi:10.1016/j.nuclphysb.2006.11.012 [arXiv:hep-ph/0609007 [hep-ph]].
- [13] P. Mastrolia and G. Ossola, “On the Integrand-Reduction Method for Two-Loop Scattering Amplitudes,” JHEP **11**, 014 (2011) doi:10.1007/JHEP11(2011)014 [arXiv:1107.6041 [hep-ph]].
- [14] S. Badger, H. Frellesvig and Y. Zhang, “Hepta-Cuts of Two-Loop Scattering Amplitudes,” JHEP **04**, 055 (2012) doi:10.1007/JHEP04(2012)055 [arXiv:1202.2019 [hep-ph]].
- [15] Y. Zhang, “Integrand-Level Reduction of Loop Amplitudes by Computational Algebraic Geometry Methods,” JHEP **09**, 042 (2012) doi:10.1007/JHEP09(2012)042 [arXiv:1205.5707 [hep-ph]].
- [16] G. Passarino and M. J. G. Veltman, “One Loop Corrections for e+ e- Annihilation Into mu+ mu- in the Weinberg Model,” Nucl. Phys. B **160**, 151-207 (1979) doi:10.1016/0550-3213(79)90234-7
- [17] K. G. Chetyrkin and F. V. Tkachov, “Integration by Parts: The Algorithm to Calculate beta Functions in 4 Loops,” Nucl. Phys. B **192**, 159-204 (1981) doi:10.1016/0550-3213(81)90199-1

- [18] F. V. Tkachov, *Phys. Lett. B* **100**, 65-68 (1981) doi:10.1016/0370-2693(81)90288-4
- [19] S. Laporta, “High precision calculation of multiloop Feynman integrals by difference equations,” *Int. J. Mod. Phys. A* **15**, 5087-5159 (2000) doi:10.1142/S0217751X00002159 [arXiv:hep-ph/0102033 [hep-ph]].
- [20] A. von Manteuffel and C. Studerus, “Reduze 2 - Distributed Feynman Integral Reduction,” [arXiv:1201.4330 [hep-ph]].
- [21] A. von Manteuffel and R. M. Schabinger, “A novel approach to integration by parts reduction,” *Phys. Lett. B* **744**, 101-104 (2015) doi:10.1016/j.physletb.2015.03.029 [arXiv:1406.4513 [hep-ph]].
- [22] P. Maierhöfer, J. Usovitsch and P. Uwer, “Kira—A Feynman integral reduction program,” *Comput. Phys. Commun.* **230**, 99-112 (2018) doi:10.1016/j.cpc.2018.04.012 [arXiv:1705.05610 [hep-ph]].
- [23] A. V. Smirnov and F. S. Chuharev, “FIRE6: Feynman Integral REduction with Modular Arithmetic,” *Comput. Phys. Commun.* **247**, 106877 (2020) doi:10.1016/j.cpc.2019.106877 [arXiv:1901.07808 [hep-ph]].
- [24] R. Britto, E. Buchbinder, F. Cachazo and B. Feng, “One-loop amplitudes of gluons in SQCD,” *Phys. Rev. D* **72**, 065012 (2005) doi:10.1103/PhysRevD.72.065012 [arXiv:hep-ph/0503132 [hep-ph]].
- [25] R. Britto, B. Feng and P. Mastrolia, “The Cut-constructible part of QCD amplitudes,” *Phys. Rev. D* **73**, 105004 (2006) doi:10.1103/PhysRevD.73.105004 [arXiv:hep-ph/0602178 [hep-ph]].
- [26] C. Anastasiou, R. Britto, B. Feng, Z. Kunszt and P. Mastrolia, “D-dimensional unitarity cut method,” *Phys. Lett. B* **645**, 213-216 (2007) doi:10.1016/j.physletb.2006.12.022 [arXiv:hep-ph/0609191 [hep-ph]].
- [27] C. Anastasiou, R. Britto, B. Feng, Z. Kunszt and P. Mastrolia, “Unitarity cuts and Reduction to master integrals in d dimensions for one-loop amplitudes,” *JHEP* **03**, 111 (2007) doi:10.1088/1126-6708/2007/03/111 [arXiv:hep-ph/0612277 [hep-ph]].
- [28] R. Britto and B. Feng, “Unitarity cuts with massive propagators and algebraic expressions for coefficients,” *Phys. Rev. D* **75**, 105006 (2007) doi:10.1103/PhysRevD.75.105006 [arXiv:hep-ph/0612089 [hep-ph]].
- [29] R. Britto and B. Feng, “Integral coefficients for one-loop amplitudes,” *JHEP* **02**, 095 (2008) doi:10.1088/1126-6708/2008/02/095 [arXiv:0711.4284 [hep-ph]].
- [30] R. Britto and E. Mirabella, “Single Cut Integration,” *JHEP* **01**, 135 (2011) doi:10.1007/JHEP01(2011)135 [arXiv:1011.2344 [hep-th]].
- [31] P. Mastrolia and S. Mizera, “Feynman Integrals and Intersection Theory,” *JHEP* **02**, 139 (2019) doi:10.1007/JHEP02(2019)139 [arXiv:1810.03818 [hep-th]].
- [32] H. Frellesvig, F. Gasparotto, M. K. Mandal, P. Mastrolia, L. Mattiazzi and S. Mizera, “Vector Space of Feynman Integrals and Multivariate Intersection Numbers,” *Phys. Rev. Lett.* **123**, no.20, 201602 (2019) doi:10.1103/PhysRevLett.123.201602 [arXiv:1907.02000 [hep-th]].
- [33] S. Mizera and A. Pokraka, “From Infinity to Four Dimensions: Higher Residue Pairings and Feynman Integrals,” *JHEP* **02**, 159 (2020) doi:10.1007/JHEP02(2020)159 [arXiv:1910.11852 [hep-th]].
- [34] H. Frellesvig, F. Gasparotto, S. Laporta, M. K. Mandal, P. Mastrolia, L. Mattiazzi and S. Mizera, “Decomposition of Feynman Integrals by Multivariate Intersection Numbers,” *JHEP* **03**, 027 (2021) doi:10.1007/JHEP03(2021)027 [arXiv:2008.04823 [hep-th]].

- [35] B. Feng, T. Li and X. Li, “Analytic tadpole coefficients of one-loop integrals,” *JHEP* **09**, 081 (2021) doi:10.1007/JHEP09(2021)081 [arXiv:2107.03744 [hep-th]].
- [36] C. Hu, T. Li and X. Li, “One-loop Feynman integral reduction by differential operators,” *Phys. Rev. D* **104**, no.11, 116014 (2021) doi:10.1103/PhysRevD.104.116014 [arXiv:2108.00772 [hep-ph]].
- [37] B. Feng, T. Li, H. Wang and Y. Zhang, “Reduction of general one-loop integrals using auxiliary vector,” *JHEP* **05**, 065 (2022) doi:10.1007/JHEP05(2022)065 [arXiv:2203.14449 [hep-th]].
- [38] B. Feng and T. Li, “PV-reduction of sunset topology with auxiliary vector,” *Commun. Theor. Phys.* **74**, no.9, 095201 (2022) doi:10.1088/1572-9494/ac7f97 [arXiv:2203.16881 [hep-ph]].
- [39] B. Feng, C. Hu, T. Li and Y. Song, “Reduction with degenerate Gram matrix for one-loop integrals,” *JHEP* **08**, 110 (2022) doi:10.1007/JHEP08(2022)110 [arXiv:2205.03000 [hep-ph]].
- [40] J. Chen and B. Feng, “Module Intersection and Uniform Formula for Iterative Reduction of One-loop Integrals,” [arXiv:2207.03767 [hep-ph]].
- [41] J. Chen, “Iteratively Reduce Auxiliary Scalar Product in Multi-loop Integrals,” [arXiv:2208.14693 [hep-ph]].
- [42] Tingfei Li, ”Non-trivial One-loop Recursive Reduction Relation”. Appearing soon.
- [43] P. A. Baikov, “Explicit solutions of the multiloop integral recurrence relations and its application,” *Nucl. Instrum. Meth. A* **389**, 347-349 (1997) doi:10.1016/S0168-9002(97)00126-5 [arXiv:hep-ph/9611449 [hep-ph]].